Origination of High T_c Copper Oxidized Superconductivity of Superconductors from Cooper Pairs Formed by Localized Electrons

Xinmin Huang^{1*} and Yu Chen²

1. School of Physics and Telecommunication Engineering, Shaanxi University of Technology, Hanzhong 723000, China

2. School of Chemistry and Environment Science, Shaanxi University of Technology, Hanzhong 723000, China

Abstract: The discovery of the superconductivity of origination of high T_c (superconducting critical temperature) copper oxidized superconductors is a significant research subject. Most of all, the origination of the electrons of Cooper pairs is not only the basis to understand the high T_c superconductive mechanism of the copper oxidized superconductors, but also is an important factor in preparing the better superconductors. In this paper, based on the theory of superconductivity of the mechanism of electronic phonons, the principle of electronegative equilibrium is applied in studying the origination of Cooper pairs which is the beginning of high T_c copper oxidized superconductivity. Combined with quantum chemical bonds theory, the conclusion is that high T_c copper oxidized superconductivity originated from localized electrons of bearing superconducting elements.

Key words: Superconductivity mechanism of phonons, principle of electronegativity equilibrium, quantum chemical bonds theory, high T_c copper oxidized superconductors, localized electrons.

1. Introduction

In 1986, Bednorz and Muller discovered that Ba-La-Cu-O oxides had the high superconducting transition temperature [1]. With the progress of the study, the researchers successively discovered more than 100 kinds of high T_c (superconducting critical temperature) superconductors of metallic elements or non-metallic elements based on copper oxides [2]. After more than 20 years' studies of theories and experiments, a comparative agreed descriptions of the high T_c copper oxidized superconductive isotopic effect and correlated experimental studies were achieved [3-8]. Now, what is clear is that the superconductivity of high T_c copper oxides originates from the mechanism of electronic phonons. In 1957, Bardeen, Cooper and Schrieffer established the micro theory of BCS (Bardeen, Cooper and Schrieffer) superconductivity based on the mechanism of the

interaction of electronic phonons [9], which put forward that the superconductive transition temperature T_c is decided by θ (θ is the inter-attraction between the electrons near Fermi interface), $\theta \sim \theta_D$ is Debye temperature and $\lambda_{eff} = N(O)V$, ($N(O)$ was a kind of spinning Bloch state density of each unit energy on Fermi interface. $V = \langle V_{kk'} \rangle_{average}$, and the energy of the transition matrix elements $V_{kk'}$ of the electronic pair in $-\hbar\omega < \varepsilon < \hbar\omega$ region is the measured average matrix element) and provided the computing formula

$$T_c = \theta e^{-\frac{1}{\lambda_{eff}}}$$
 . In high T_c copper oxidized superconductors, transition of electrons occurred between elements because of the bonding between metallic elements and oxygenic elements. Metals lost electrons while non-metal oxygen got electrons. The electrons are confixed between the lattice and the movable electrons are very few. The normal state is insulator and the distribution of the electronic states is completely different from that of metallic

Corresponding author: Xinmin Huang, professor, research fields: superconductivity. E-mail: wangjianwu718@163.com.

superconductors. Thus, the electronic pair origination of the high T_c copper oxidized became an important issue. The electronegative equilibrium principle is adopted to discuss the nature of the chemical bonds of the high T_c copper oxidized superconductors and obtained significant conclusions.

2. Methods

Sanderson pointed out [10] that in the process of forming the materials of multi electronic system by multi atoms, the difference of electro negativity of the elements in every part in the system caused electrons to transfer from the low electro negativity region to the high electro negativity region (that is, from the high chemical potential region to the low chemical potential region) to bring the electro negativity of each element in the materials equation forming electronic system to balance.

$$\text{That is: } x_a = x_b = \dots = X \quad (1)$$

x_a, x_b represent the electro negativity of different elements.

In addition, there is a condition of charge:

$$\sum_a q_a = Q \quad (2)$$

X and Q are the electro negativity and total charge, respectively.

In the superconductors bonded formed between the elements, heterogeneity formed because of the existences of the below states.

(1)The atomic natures which formed superconductors and the distributions are different.

(2)The atoms which formed superconductors were same but the bonding ways and distribution are different.

(3)The atoms which formed superconductors and the bonding ways are same but the types of lattice are different.

According to the electronegativities principle in different regions, it could be obtained:

$$x_I = x_{II} = \dots = x \quad (3)$$

$x_I, x_{II} \dots$ are regional electronegativities in different regions.

In the same region:

$$x_a = x_b = \dots = x_I \quad (4)$$

$$x'_a = x'_b = \dots = x_{II} \quad (5)$$

$$x^n_a = x^n_b = \dots = x_n \quad (6)$$

$x_a, x'_a, \dots, x^n_a, x_b, x'_b, \dots, x^n_b$ are the electronegativities of the elements in different regions.

But among $x_a, x'_a, \dots, x^n_a, x_b, x'_b, \dots, x^n_b$, they were not necessarily equal.

In the structure of the high T_c copper oxidized superconductors, according to the different electric properties, the lattices of the high T_c copper oxidized superconductors are divided into conducting layer and carrier layer while the same lattice had:

$$x_{\text{conductivity layer}} = x_{\text{carrier reservoir}} \quad [11].$$

3. Characteristics of the Elements Bonds and Superconductivity in the High T_c Copper Oxidized Superconductors

3.1 Characteristics of Elements Bonding

Pauling [12] holds the opinion that the iconicity of bonds is caused by the differences in the capabilities of electronic attraction of the atoms in molecules. When the attraction capability of electrons by atom B is bigger than atom A and it is big enough to make it form A^+B^- , this bond is expressed as the ideal ionic bonds. Then, the theoretical value of dipole moment was $\mu_0 = eR$. When the attraction capability of electrons by atom B is not big enough to form A^+B^- and the charge of bonded electrons slightly deviated to atom B, some ionic bonds are formed.

Hooydonk [13] gave a simple computing method of calculating the relationship between the high iconicity and electronegativity of the bonds.

$$v\% = \left| \frac{X_A - X_B}{X_A + X_B} \right| \times 100\% \quad (7)$$

Table 1 The $-\delta_0$ value of the high T_c copper oxidized superconductors and T_c reported in literature.

NO	Superconductor	$-\delta_0$	T_c / K
1	La _{1.85} Ba _{0.15} CuO _{3.93+x}	0.494	26
2	La _{1.85} Sr _{0.15} CuO _{3.93+x}	0.491	37
3	LaBa ₂ Cu ₃ O _{6.5+x}	0.455	75
4	Y _{0.5} La _{0.5} Ba ₂ Cu ₃ O _{6.5+x}	0.451	82
5	YBa ₂ Cu ₃ O _{6.5+x}	0.447	94
6	Ti ₂ Ca ₂ Ba ₂ Cu ₃ O _{10+x}	0.395	125
7	Bi _{1.6} Pb _{0.3} Sb _{0.1} Sr ₂ Ca ₂ Cu ₃ O _{9.85+x}	0.386	132

$v\%$ represents the percentage of ionicity. X_A , X_B are the electroneativity of atom A and atom B.

The value of the ionicity of the bonds obtained by Hooydonk's method and by Matcha's [13] are nearly consistent.

3.2 Characteristics of Elements Bonding in the High T_c Copper Oxidized Superconductors

Zhao Liangzhong [14] studied the Cu-O bond types of the high T_c copper oxidized superconductors and some charges. The results were listed in Table 1.

After Liu Honglin and Chen Nianyi [15] conducted the ab initio of full-electronic computation of quantum chemistry in YBa₂B₃O₇ atomic cluster autonomous crystalline field, they believed that in YBa₂B₃O₇, there existed $[\text{CuO}_2\text{-O-CuO-O-CuO}_2]^{-6.8}$ complex ions, which are conjoint with covalent bonds in the single cell while they were conjoint with ionic bonds between the single cells in the direction of Axis C^[16-20].

4. Conclusions

On the basis of the superconductive theory of the electronic phonon mechanism, the principle of electronegativity equilibrium was adopted to study the electronic origination electronic pairs originated from the high T_c copper oxidized superconductivity. It was proposed that the high T_c copper oxidized superconductivity originated from the mechanism of electronic phonons and the electrons forming Cooper pairs were those influenced by different interactions of elements Cu and O on CuO₂ plane to make the

localized electros which corresponded to $V_{Cu} \approx V_o$.

References

- [1] Bednorz, J. G., Muller, K. A. 1986. "Possible T_c Superconductivity in the Ba-La-Cu-O System." *Z. Phy. B* 64: 189-193.
- [2] Zhou, W. Z., and Liang, W. Y. 1999. *Research on High T_c Superconductive Basis*. Shanghai: Shanghai Scientific and Technological Press.
- [3] Frank, J. P., Harker, S., and Brewer, J. H. 1993. "Copper and Oxygen Isotope Effects in La_{2-x}Sr_xCuO₄." *Phys. Rev. Lett.* 71: 283-6.
- [4] Chen, Y., Huang, X. M., and Wang, J. W. 2008. "Relationship between Isotopic Effects of High T_c Copper Oxidized Superconductors and Contents of Oxygen and Position." *Journal of Low T_c Physics* 30 (2): 158-160.
- [5] Chen, Y., Huang, X. M., and Wang, J. W. 2012. "Correlcotions between the Isotope Effects of the Cu and O Elements in High T_c Copper Oxide Superconductors." *Journal of Materials and Engineering B* 2: 404-7.
- [6] Vladimir, E., Manuchargan, J. K., Glazman, L. I., and Devoret, M. H. 2009. "Fluxonium: Single Cooper Pair Circuit Free of Charge Offset." *Science* 326 (2): 113-6.
- [7] Chen, Y., Huang, X. M., and Wang, J. W. 2011. "The Interactions between Electrons and Phonons in Bonded Elements." *Journal of Materials and Engineering B* 1: 227-231.
- [8] Logvenov, G., Gozar, A., and Bozovic, I. 2009. "High Temperature Superconductivity in a Single Copper Oxygen Plane." *Science* 236 (30): 699-702.
- [9] Bardeel, J., Cooper, L. N., and Schrieffer, J. R. 1957. "Theory of Superconductivity." *Phys. Review* 108: 1175-1204.
- [10] Sanderson, R. T. 1976. *Chemical bond and energy (Second Edition)*. Washington: Academic Press, 126-130.
- [11] Chen, Y., and Huang, X. M. 2009. "Some Superconductive Influential Characteristics of 1- ω in Superconductor." *Journal of Low T_c Physics* 31 (1): 80-82.
- [12] Pauling, L. 1960. *The Nature of Chemical Bond and the Structure of Molecules and Crystals*. California: Corne University Press.
- [13] Yang, P. 2007. "Parameters of Molecular Structure and its Relevant Law of Physical Property." Beijing: Science Press.
- [14] Zhao, L. Z. 1995. "Cu-O Bond Types and Some Charges of Oxygen in High T_c Superconductor." *Journal of Physical Chemistry* 11 (1): 1051-7.
- [15] Liu, H. L., and Chen, N. Y. 1992. "Ab Initio of Full-Electronic Computation of Quantum chemistry in YBa₂B₃O₇ Atomic Cluster Autonomous Crystalline

- Field.” *Chinese Science (B)* 25 (2): 132-6.
- [16] Chen, Y., Huang, X. M., and Wang, J. W. 2012. “Correlation Between the Isotope Effects of the Cu and O Elements in High T_c Copper Oxide Superconductors.” *Journal of Materials Science and Engineering B* 2: 404-7.
- [17] Chen, Y., Li Y., and Yao. J. B. 2014. “The Characteristics of the Cooper Pair Electrons in High T_c Copper Oxide Superconductors.” *Journal of Materials Science and Engineering A* 4 (5): 160-3.
- [18] Huang, X. M., and Chen, Y. 2014. “Characteristics of the Origin of High T_c Copper Oxidized Copper Pairs.” *Journal of Materials Science and Engineering B* 4 (4): 129-133.
- [19] Huang, X. M., and Chen, Y. 2014. “Two Subsystems and Chemical Bonds in High Copper Oxidized Superconductors.” *Journal of Materials Science and Engineering A* 4 (7): 255-8.
- [20] Chen, Y., Yao, J. B., and Yu, L. 2014. “Characteristics of Superconductivity in Superconductors.” *Journal of Materials Science and Engineering B* 4 (6): 198-202.