Seasonal Differences in Live Foraminiferal Densities: Case Studies from Tropical and Temperate Intertidal Environments

Ashleigh Costelloe and Brent Wilson

Department of Chemical Engineering, The University of the West Indies, St. Augustine, Trinidad

Abstract: Statistically significant differences (when $p < 0.05$ using the permutation t-test) among Live Foraminiferal Densities (LFDs) recovered in August, November, March and May were detected at tropical Caroni swamp, Claxton bay (Trinidad), temperate Cowpen marsh and Brancaster marsh (U.K.). The monthly mean LFDs of the foraminiferal metacommunities (all stations), assemblages (groups of stations defined by cluster analysis), and the agglutinated and calcareous specimens within each were compared separately. The LFDs of the Caroni swamp metacommunity did not fluctuate significantly among months, but significantly higher abundances of agglutinated specimens occurred in the upper assemblage in March; and calcareous specimens in the lower assemblage in November. At Claxton bay, monthly LFDs of the metacommunities and assemblages did not vary significantly, but calcareous and agglutinated species within each favoured dry (March and May) and wet (August and November) months respectively. At the temperate marshes, significantly higher LFDs of the metacommunities and assemblages were recorded in warmer months. August blooms of the Cowpen marsh metacommunity was attributed to agglutinated specimens in the upper assemblage, and calcareous specimens in the lower assemblage. May blooms of the Brancaster marsh upper assemblage were attributed to calcareous specimens, but there were no seasonal blooms of the lower assemblage.

Key words: Foraminifera, marsh, swamp, intertidal, seasonal, trends, densities.

1. Introduction

Foraminifera are used for environmental monitoring within marginal marine habitats, where low foraminiferal densities have been suggested to indicate anthropogenic stress [1-7]. However, fluctuations in foraminiferal densities could be linked to seasonal changes, low densities occurring during unfavourable seasons [8, 9]. This paper investigates seasonality of foraminiferal population dynamics from tropical swamps and temperate marshes.

In the tropics, seasonal shifts of the inter-tropical convergence zone cause rainfall to vary more dramatically than the average temperature. The seasonal impact on the tropical intertidal foraminiferal community is only vaguely understood [10, 11]. The

temperate seasonal trend is better understood, spring or summer blooms in foraminiferal populations being associated with increased food supply [12-14].

Live Foraminiferal Densities (LFDs) from two tropical mangal swamps—Caroni swamp and Claxton bay (Fig. 1) were compared to two temperate marshes—Cowpen marsh and Brancaster marsh. The LFDs were compared among March, May, August and November. At the tropical swamps, March and May represent early and late dry season, and August and November represent early and late wet season. At the temperate locations, conditions during early and late spring were represented by March and May, summer by August, and late autumn by November.

The vertical distribution of intertidal foraminifera is closely linked to elevation relative to the tidal frame [15]. Increased aerial exposure above mean high water results in increased dissolved oxygen and salinity (the latter being due to evaporation). Foraminiferal

Corresponding author: Ashleigh Costelloe, Ph.D., main research fields: environmental micropaleontology and biostratigraphy.

Fig. 1 Map of tropical study locations in Trinidad, SE Caribbean (a), Caroni swamp (b), where three sites were sampled (c), Claxton bay (d) where one site was sampled (e).

assemblages at higher elevations are almost exclusively agglutinated while calcareous species dominate at lower elevations [16]. At each location, monthly LFDs of (a) metacommunities (all stations), (b) assemblages (defined by cluster analysis) and (c) calcareous and agglutinated specimens within each were compared.

Foraminiferal assemblages vary between months, yet one assemblage scheme must be applied to all months for comparison of temporal variations. The dead or total (dead + live) counts limit some temporal variability by incorporating dead tests from previous months. Thus, seasonal perturbations are removed, giving an average annual representation of foraminiferal distributions for defining one assemblage scheme [17]. Therefore, either dead or total counts were used for defining assemblages, although this paper examines the live foraminiferal community among months. At Caroni swamp, an assemblage scheme constrained by elevation was predefined by Wilson et al. [18]. The samples recovered from Caroni swamp and Claxton bay in this study were combined with those of Wilson et al. [18] in separate cluster analyses. This was done to validate the cluster analyses where few (< 15) stations were sampled.

2. Locations of Case Studies

Caroni swamp (10°27'38" N, 61°14'55" W) and Claxton bay (10°21'0" N, 61°38'0" W) are 35 km apart, on the west coast of Trinidad. They open to the Gulf of Paria—a semi-enclosed embayment separating Trinidad from Venezuela (Fig. 1a). Salinity in the Gulf of Paria is consistently lower than that of tropical Atlantic surface water [19] due to freshwater input from the Orinoco river in Venezuela, which has maximum discharge rates from September to November [20]. Caroni swamp is a 60 km² mangrove wetland reserve (Fig. 1b). It is micro-tidal with semi-diurnal tides and a spring tidal range of about 1.2 m [18]. Claxton bay is a 2.25 km stretch of mangrove

shoreline in an industrial area with heavy shipping activity (Fig. 1c). In Trinidad, the months of January to May traditionally receives heavier rainfall than June to December which is drier (Fig. 2).

Cowpen marsh (54°37'18" N, 1°13'29" W) is a 400 m wide tidal flat located on the western side of the Tees estuary, NE England (Figs. 3a and 3b). It exhibits a macro-tidal range between 6.0 m (spring tide) and 4.6 m (neap tide) [21]. Brancaster marsh (52°57'44" N, 0°38'29" E) is approximately 150 m wide, located on the eastern Norfolk coast of England (Figs. 3a and 3c), approximately 300 km south of Cowpen marsh. Brancaster marsh exhibits a macro tidal range between 6.0 m (spring tide) and 3.1 m (neap tide) [22]. Seasonal averages of maximum and minimum temperatures and rainfall at Boulmer (32 km north of Cowpen marsh) and Weybourne (33 km south of Brancaster) are shown in Fig. 4.

3. Materials and Methods

3.1 Sample Collection and Preparation

At Caroni swamp and Claxton bay, four replicate sediment samples (75 cm² × 1 cm) were collected twice in the wet (August and November 2012) and dry (March and May 2013) seasons. Three sites at Caroni swamp (Fig. 1d) and one site at Claxton bay (Fig. 1e) were sampled at high (stations a), mid (stations b) and low (stations c) tidal elevations, which was determined

Fig. 2 Trends in the 2011-2013 rainfall recorded by the Trinidad and Tobago Meteorological Services located at the Piarco International Airport (13.5 km east of Caroni swamp and 27.5 km northeast of Claxton bay).

Fig. 3 Map of temperate study location in eastern England, UK (a): Cowpen Marsh where one site was sampled (b) and Brancaster Marsh where one site was sampled (c).

Fig. 4 The average monthly maximum and minimum temperatures (a) and rainfall (b) at each of the closest coastal met office locations to Cowpen (Boulmer, UK) and Brancaster (Weybourne, UK) based on historical data from 1981 to 2010 recorded by the UK meteorological offices.

by surveying (Table 1). The transects from stations a to c were 7 m, 12 m, and 15 m-long at Caroni swamp sites 1-3 respectively, and 23 m-long at Claxton bay.

Samples were soaked in a solution of the protein stain rose Bengal and 70% isopropyl alcohol for 15 days to stain the cytoplasm of live specimens [23].

Samples were then washed through 2,000 μm and 63 μm sieves. The barren > 2,000 μm fraction was discarded. Foraminifera from Caroni swamp were picked wet and counted from the entire 63-2,000 μm fraction. Samples from Claxton bay were split into 1/16 aliquots using a wet microsplitter and used as representative subsamples [24]. The taxonomy

follows Ruth Todd and Paul Bronnimann [25].

At Cowpen marsh, 31 stations were sampled along a 200 m-long transect on 1 May 1995 (late spring), 10 August 1995 (summer), 7 November 1995 (autumn) and 5 March 1996 (late spring). At Brancaster marsh, 24 stations were sampled along 94 m-long transect on 24 November 1995 (autumn), 1 March 1996 (early spring), 25 May 1996 (late spring) and 16 August 1996 (summer). The elevations of the stations were determined by surveying (Table 2). One sediment sample ($10 \text{ cm}^2 \times 1 \text{ cm}$) was collected from each station at both locations and soaked in ethanol or formalin with rose Bengal. Each sample was wet sieved through 500 μm and 63 μm sieves and the > 500 μm fraction was discarded. Foraminiferans 63-500 μm were counted wet from 1/8 aliquots of each sample using a wet splitter. The taxonomy follows

John Murray [26, 27] and Sachade Rijk [28]. Minor species were grouped at the genus level (e.g. *Quinqueloculina* spp.) and the combination of possible species or echophenotypes were grouped under a single species name (*Ammonia* spp.).

3.2 Foraminiferal Assemblages

Cluster analysis used total counts from Caroni swamp and Claxton bay to emulate methods used for an assemblage scheme predefined by Brent Wilson et al. [18]. The dead counts from Cowpen marsh and Brancaster marsh were used since an analysis by Benjamin Horton and John Murray [21] to determined dead counts to better represent average annual conditions.

The recovery from replicated samples collected at Caroni swamp and Claxton bay was summed to

Table 1 Elevations of stations sampled at Caroni swamp and Claxton bay.

Tropical study locations	Elevation (above mean sea level (m))		
Caroni swamp	Station a	Station b	Station c
Site 1	0.655	0.453	-0.176
Site 2	1.487	0.195	-0.051
Site 3	-0.14	-0.102	-0.117
Claxton bay	Station a	Station b	Station c
Site 1	0.228	0.051	-0.151

Table 2 Elevations of stations sampled at Cowpen marsh and Brancaster marsh.

Cowpen marsh				Brancaster marsh			
Station	Elevation (m, OD)	Station	Elevation (m, OD)	Station	Elevation (m, OD)	Station	Elevation (m, OD)
1	3.24	17	1.99	1	3.27	17	2.98
2	2.86	18	1.94	2	3.21	18	3.07
3	2.81	19	1.88	3	3.24	19	3.02
4	2.57	20	1.74	4	3.23	20	3
5	2.47	21	1.52	5	3.21	21	2.48
6	2.42	22	1.32	6	3.22	22	2.31
7	2.38	23	1.13	7	3.22	23	1.88
8	2.33	24	0.93	8	3.23	24	1.85
9	2.27	25	0.72	9	3.22		
10	2.22	26	0.54	10	3.24		
11	2.18	27	0.36	11	N/A		
12	2.22	28	0.17	12	3.21		
13	2.17	29	-0.04	13	3.19		
14	2.14	30	-0.22	14	3.19		
15	2.09	31	-0.35	15	3.14		
16	2.04			16	3.03		

overcome sampling errors caused by the patchy distribution of foraminiferans [29]. In separate analyses, the stations from Caroni swamp and Claxton bay sampled in May 2012 and 2013 respectively were clustered with the stations of Brent Wilson et al. [18] sampled in May 2006. The month with the highest density recovered from Cowpen marsh (August 10, 1995) and Brancaster marsh (November 24, 1995) was used for cluster analysis. Q-mode cluster analysis in PAST 3.0 was used, with the correlation coefficient and the unweighted pair-group average linkage method [31]. Stations where counts were < 50 specimens and taxa with a recovery $< 2\%$ of the total recovery were omitted.

3.3 Comparison of Live Foraminiferal Densities

The monthly LFDs recorded are a function of the number of samples and volume of sediment collected at each location. To facilitate comparisons between locations, the mean LFD (sum of specimens/number of samples) was used. The foraminiferal counts from Cowpen marsh and Brancaster marsh were normalized to 75 cm^3 of sediment. Statistical comparison of mean LFDs used the permutation t-test in PAST 3.0 [30].

The different data sets had a non-normal distribution with unequal variances. The number of samples per site were small ($N < 50$) and differed between locations. When using small data sets with non-normal distributions, the permutation t-test gives more accurate results than a two sample t-test or one way ANOVA [31]. The observed t statistic (the normalized difference between means) was compared with the t statistics from 1,000 random pairs of replicates from the pooled data set. The permutation t-test gives the probability (p) that the means of the data sets are different within a 95% confidence interval ($\alpha = 0.05$). Only the p-value and a reliable confidence interval are necessary to determine statistical significance. Where p was < 0.05 , the two data sets had significantly different means.

4. Results

4.1 Caroni Swamp

There was no significant difference ($p > 0.05$ for all comparisons) between mean LFDs of the foraminiferal metacommunity recorded at Caroni swamp in August, November, March or May (Fig. 5a). Agglutinated specimens dominated except in November when calcareous specimens comprised 62% of the metacommunity (Fig. 5b). In August, *Ammonium salsum* (32%) and *Trochammina advena* (22%) co-dominated, with fewer *Arenoparella mexicana* (14%) and *Miliammina fusca* (11%). In November, *Ammonia beccarii* (49%) dominated with fewer *A. salsum* (22%) and *Ammonia tepida* (13%). In March, *M. fusca* (29%)

Fig. 5 Mean LFDs of the Caroni Swamp metacommunity (a), calcareous (b) and agglutinated (c) specimens within.

dominated with fewer *A. salsum* (19%) and *T. advena* (14%). In May, *A. beccarii* (48%) was dominant, with fewer *T. advena* (21%) and *A. salsum* (10%). A significant increase in the number of calcareous specimens (mainly *A. beccarii*) occurred in November (Fig. 5c).

Cluster analysis divided the Caroni swamp metacommunity into upper (stations 1a, 1c and 2a) and mid (1b, 2b, 2c, 3a, 3b and 3c) assemblages, corresponding to zones I and II, respectively, established by Brent Wilson et al. [18]. A lower assemblage (zone III) was not represented at Caroni swamp in this study, and the foraminiferal assemblages were not constrained by elevation.

The mean LFD of the upper assemblage was significantly greater in March than all other months (Fig. 6a). Agglutinated specimens dominated (> 80%), except in November (47%). In August, *Arenoparella mexicana* (43%) dominated, with fewer *Trochammina inflata* (33%) and *A. salsum* (11%). In November, *A. beccarii* (44%) dominated, with fewer *A. salsum* (13%) and *A. mexicana* (10%). In March, *M. fusca* (31%) dominated, with fewer *A. salsum* (18%) and *T. advena* (13%). In May, *T. inflata* (34%) dominated, with fewer *A. mexicana* (19%), *T. advena* (13%) and *A. beccarii* (13%). Calcareous specimens comprised 1% of the assemblage in August which was significantly lower than all other months (Fig. 6b). The highest abundance of agglutinated specimens, mostly *M. fusca*, *A. salsum* and *T. advena* were recorded in March, which were significantly greater than in November and May (August just failed to meet the criteria for significant difference, Fig. 6c).

The mean LFD of the mid assemblage recorded in November was significantly greater than all other months (Fig. 7a). The mid foraminiferal assemblage was dominated by agglutinated specimens in August (97%) and March (72%, Fig. 7b), and by calcareous specimens in November (63%) and May (53%, Fig. 7c). In August and March, *A. salsum* (38% and 20%, respectively), *M. fusca* (13% and 26%, respectively)

Fig. 6 Mean LFDs of the Caroni Swamp upper assemblage (a), calcareous (b) and agglutinated (c) specimens within.

and *T. advena* (26% and 18%, respectively) were abundant. In March, *A. tepida* (17%) and *A. beccarii* (11%) were abundant. In November and May, *A. beccarii* (49% and 48%, respectively) dominated. In November, *A. salsum* (20%) and *A. tepida* (13%) and in May, *T. advena* (23%) and *A. salsum* (10%) were abundant. The abundance of calcareous specimens (mostly *A. beccarii*) fluctuated significantly among all months, but was ≥ 3 times greater in November (Fig. 7c).

4.2 Claxton Bay

The mean LFDs of the metacommunity at Claxton bay did not differ significantly among months (Fig. 8a).

Seasonal Differences in Live Foraminiferal Densities: Case Studies from Tropical and Temperate Intertidal Environments

Fig. 7 Mean LFDs of the Caroni Swamp mid assemblage (a), calcareous (b) and agglutinated (c) specimens within.

Agglutinated specimens were significantly greater in November than March and May (Fig. 8b). There was also a significant decrease in the abundance of calcareous specimens recorded in November compared to March and May (Fig. 8c). During the wet months of August and November, *T. advena* dominated (69% and 66%, respectively), with fewer *A. tepida* (12% and 15%, respectively) and *A. salsum* (15%) in November only. In the dry months of March and May, *T. advena* was also dominant (50% and 43%, respectively), but *A. tepida* (27% and 30%, respectively) was more abundant than in the wet months. Conditions at Claxton bay in the wet season were unfavourable to the population growth of calcareous specimens, which achieved higher abundances in the dry season.

Fig. 8 Mean LFDs of the Claxton Bay metacommunity (a), calcareous (b) and agglutinated (c) specimens within.

The Claxton bay metacommunity was divided into mid (stations 1b) and lower (stations 1a and 1c) assemblages, relative to Brent Wilson et al. [18] zones II and III, respectively. An upper assemblage (zone I) was not represented at Claxton bay in this study. Foraminiferal assemblages were not constrained by elevation.

The mean LFDs of the mid assemblage recorded at Claxton bay were statistically similar among months (Fig. 9a). The abundance of agglutinated specimens was significantly greatest in August (Fig. 9b). Monthly mean abundances of calcareous specimens were similar among months (Fig. 9c). *Trochammina advena* always dominated, forming 80% in August, 66% in November, 71% in March, and 56% in May.

Fig. 9 Mean LFDs of the Claxton bay mid assemblage (a), calcareous (b) and agglutinated (c) specimens within.

In November and May, *A. tepida* (13% and 14%, respectively) and *A. salsum* (16% and 12%, respectively) were also abundant. Wet conditions in August favoured population growth of agglutinated specimens, mainly *T. advena*.

The monthly mean LFDs recorded for the lower assemblage at Claxton bay were not significantly different (Fig. 10a). Calcareous specimens were significantly fewer and agglutinated specimens denser in the late wet season (November) than in the dry season (March and May, Figs. 10b and 10c). In August and November (wet months), *T. advena* dominated (55% and 66%, respectively), with fewer *A. tepida* (17%

Fig. 10 Mean LFDs of the Claxton Baylower assemblage (a), calcareous (b) and agglutinated (c) specimens within.

and 16%, respectively). In August and November, *Triloculina oblonga* (19%) *A. salsum* (15%) were abundant, respectively. In March and May (dry months), *A. tepida* (38% and 41%, respectively) and *T. advena* (37% and 33%, respectively) co-dominated, with fewer *T. oblonga* (11%) in May only. Favourable population growth of agglutinated species in the lower assemblage occurred during wet conditions and calcareous specimens during dry conditions.

4.3 Cowpen Marsh

The mean LFD of the foraminiferal metacommunity was significantly greatest in August at Cowpen marsh

(Fig. 11a). The monthly abundances of calcareous specimens did not differ significantly (Fig. 11b), but the abundance of agglutinated specimens recorded in March was significantly less when compared to August only (Fig. 11c). *Jadammina macrescens* always dominated the foraminiferal metacommunity: 47% in May, 37% in August, 40% in November, and 29% in March. *Haynesina germanica* was always abundant, but more so in August and March (24% and 23%, respectively) than May and November (15% and 14%, respectively). *Trochammina inflata* was abundant in May, August and November (11%, 15% and 19%, respectively) and *Quinqueloculina* spp. (11%) was abundant in November only. In March, *Quinqueloculina*

Fig. 11 Mean LFDs of the Cowpen Marsh metacommunity (a), calcareous (b) and agglutinated (c) specimens within.

spp. (17%) and *Elphidium williamsoni* (14%) were abundant. The monthly mean abundance of agglutinated specimens (mainly *J. macrescens*) increased from late spring to summer, when a significant maximum was recorded in comparison to early spring.

Cluster analysis divided the foraminiferal metacommunity at Cowpen marsh into an upper assemblage (stations 1-16) between 3.24 m and 2.04 m above Ordnance Datum (OD), and a lower assemblage (stations 17-31) between 1.99 m and -0.04 m above OD.

The mean LFD of the upper assemblage was significantly greater in August than May and March, but not November (Fig. 12a). *Jadammina macrescens* always dominated: 64% in May, 58% in August, 47% in November and 37% in March. In May, August and November, *T. inflata* (15%, 24% and 22%, respectively) was abundant. In November and March, *Quinqueloculina* spp. (13% and 20%, respectively) was abundant, and *H. germanica* (18%) in March only. A summer (August) peak in agglutinated specimens (mainly *J. macrescens* and *T. inflata*) existed, with values declining in November, but only achieving a significant minimum in May (Fig. 12b). At that time, calcareous specimens (mainly *H. germanica*) significantly increased in comparison to the warmer months of May and August (Fig. 12c).

The mean LFD of the lower assemblage was significantly greater in August than in March and November (Fig. 13a). The number of calcareous specimens was significantly highest in August when compared to November and March but not May (Fig. 13b). Similarly low abundances of agglutinated specimens were recorded for all months, except November, which had zero variance such that the permutation t-test could not be performed (Fig. 13c). *Haynesina germanica* dominated in May (44%), August (52%), November (64%) and March (39%). *Quinqueloculina* spp. was subdominant in May (23%) and August (22%), but *E. williamsoni* was subdominant in November (24%) and March (24%). Calcareous specimens (mainly *H. germanica* and

Fig. 12 Mean LFDs of the Cowpen Marshupper assemblage (a), calcareous (b) and agglutinated (c) specimens within.

Quinqueloculina spp.) increased in May to achieve a significant maximum in August and minima in November and March.

4.4 Brancaster Marsh

The mean LFD of the foraminiferal metacommunity at Brancaster marsh was significantly highest in May (Fig. 14a). In May and August, the metacommunity was dominated by *Quinqueloculina* spp. (28% and 46%, respectively), with fewer *J. macrescens* (16% and 17%, respectively) and *T. inflata* (15% and 12%, respectively). In November and March, *J. macrescens* (35% and 27%, respectively) and *T. inflata* (23% and 26%, respectively) co-dominated with fewer *Quinqueloculina*

Fig. 13 Mean LFDs of the Cowpen Marshlower assemblage (a), calcareous (b) and agglutinated (c) specimens within.

spp. (17% and 15%, respectively). *Haynesina germanica* was only abundant in March (16%) and *Cyclogyra involvens* in August (11%). The abundance of calcareous specimens (mostly *Quinqueloculina* spp.) was significantly higher in May compared to November and March, and lower in November compared to August and May (Fig. 14b). The abundance of agglutinated specimens significantly decreased in August compared to November and May (Fig. 14c).

Cluster analysis revealed two foraminiferal assemblages constrained by elevation. The upper assemblage comprised stations 1 to 19 (3.27-3.02 m OD). The lower assemblage comprised stations 20 to 24 (3.0-1.85 m OD).

Seasonal Differences in Live Foraminiferal Densities: Case Studies from Tropical and Temperate Intertidal Environments

Fig. 14 Mean LFDs of the Brancaster Marshmetacommunity (a), calcareous (b) and agglutinated (c) specimens within.

The mean LFD of the upper assemblage was significantly highest in May (Fig. 15a). In November and March, *J. macrescens* (32% and 31%, respectively) and *T. inflata* (27% and 31%, respectively) co-dominated, with fewer *Quinqueloculina* spp. (16% in both months). In May and August, *Quinqueloculina* spp. (29% and 41%, respectively) dominated with fewer *J. macrescens* (17% and 20%, respectively) and *T. inflata* (17% and 16%, respectively). A significant influx of calcareous specimens occurred in May (Fig. 15b), and a significant decline in agglutinated specimens occurred in August (Fig. 15c). Therefore, the late spring bloom in the upper assemblage could be attributed to an increase of *Quinqueloculina* spp.

Fig. 15 Mean LFDs of the Brancaster Marshupper assemblage (a), calcareous (b) and agglutinated (c) specimens within.

and *E. williamsoni*, which were the only abundant calcareous species in May.

There was no significant difference between monthly mean LFDs (Fig. 16a), or the abundance of calcareous (Fig. 16b) and agglutinated (Fig. 16c) specimens recorded for the lower assemblage at Brancaster marsh. Calcareous specimens dominated, except in November, when *J. macrescens* (52%) dominated with fewer *Quinqueloculina* spp. (24%) and *H. germanica* (16%). In March and May, *H. germanica* (67% and 60%, respectively) was dominant, with fewer *Quinqueloculina* spp. (11% and 15%, respectively). In August, *Quinqueloculina* spp. (63%) was dominant, with fewer *H. germanica* (24%) and *J. macrescens* (10%).

Fig. 16 Mean LFDs of the Brancaster Marsh assemblage (a), calcareous (b) and agglutinated (c) specimens within.

5. Discussion

At Caroni swamp, the upper swamp LFD increased in the early dry season (March), led by the agglutinated foraminiferal population (*M. fusca*, *A. salsum* and *T. advena*). The mid swamp LFD increased in the late wet season (November), mainly through calcareous specimens (*A. beccarii*). The mean LFDs of the metacommunity and assemblages at Claxton bay were consistent among months, but there were seasonal patterns in the abundances of agglutinated and calcareous specimens within. At Claxton bay, agglutinated specimens were more abundant in the wet season, while calcareous

specimens were more abundant in the dry season. Population growth of agglutinated and calcareous specimens at Caroni swamp did not follow the same seasonal pattern as at Claxton bay.

The population dynamics of selected foraminiferal species in Jamaica were described as aseasonal [10]. Abiotic factors, which are more influential in temperate regions, were considered secondary to the biotic controls in tropical habitats. Elsewhere in the Caribbean region, seasonal population dynamics have been observed. The mean diversities of the epiphytal foraminiferal communities in Nevis were lowest during a dry season and abundances increased after a hurricane season [9]. In the Cariaco basin (Venezuela), Orinoco river discharge rates vary seasonally, causing variations in the thermocline and chlorophyll maximum which produce seasonal variations in planktonic foraminifera [32]. At Kaw estuary (French Guiana), calcareous specimens suffered dissolution due to reduced pH levels during the wet season. Increased fluvial runoff from the Amazon river lowered salinity and increased deposition of muds, which reduced pH levels [33]. Trinidad is influenced similarly by the Orinoco river plume during the wet season. The foraminiferal assemblages at Claxton bay exhibited seasonal changes analogous to those of Kaw estuary [33], but not at Caroni swamp.

At the temperate marshes, LFDs were highest in the warmer months: August at Cowpen marsh and May at Brancaster marsh. The upper Cowpen marsh was populated by agglutinated species (*J. macrescens* and *T. inflata*), which bloomed in the summer. During the unfavourable cooler month of March, *H. germanica* significantly increased. In the lower marsh, *H. germanica* was always dominant, and bloomed with other calcareous species (mostly *Quinqueloculina* spp. and *E. williamsoni*) in the summer. At Brancaster marsh, late spring blooms in the upper marsh were attributed to the growth of *Quinqueloculina* spp. and *E. williamsoni*. No significant fluctuations were observed for the lower Brancaster marsh.

Spring and summer blooms in temperate foraminiferal communities are common and have been related to food supply [12, 14, 28, 34-39]. Phytoplankton blooms in the summer and spring are mainly an offshore occurrence, but can be transported onshore by tides [40]. Additionally, spring and summer algal blooms have been known to occur in the marsh sediment, which may promote growth of foraminiferal communities feeding on them [12, 13, 28].

Drawing a general conclusion for a region based on a single study location is not possible. The patterns observed at one location are the result of unique environmental characteristics. The geography and vegetation of an area can greatly influence the abiotic and biotic variables that regulate foraminiferal populations.

6. Conclusion

At Caroni swamp, dry months were more favorable for foraminiferal population growth of the upper assemblage ((mainly *M. fusca*, *A. salsum* and *T. advena*). Population growth of the mid assemblage (mainly *A. beccarii*) was favored under wet conditions. At Claxton bay, population growth of calcareous (mainly *A. tepida* and *T. oblonga*) and agglutinated (mainly *T. advena*) specimens favoured dry and wet months, respectively.

At the temperate marshes, higher mean LFDs were recorded in warmer months. Summer blooms of the Cowpen marsh upper assemblage was attributed to agglutinated specimens (mainly *J. macrescens* and *T. inflata*) and calcareous specimens (mainly *H. germanica*) in the lower assemblage. Spring blooms of the Brancaster marsh upper assemblage was attributed to calcareous specimens (mainly *Quinqueloculina* spp. and *E. williamsoni*), but there were no seasonal blooms in LFDs of the lower assemblage.

The seasonal trend at one environment may not be the same at another, even though both environments are in close proximity and within the same climatic zone. Seasonal trends include significant changes in

foraminiferal densities and species compositions. It is important to detect natural fluctuations in foraminiferal population dynamics of a study area before assessing anthropogenic impacts on foraminiferal communities.

Acknowledgements

This research was supported by the Campus Research and Publication Fund to the Geoscience Unit at the University of the West Indies (St. Augustine).

References

- [1] Alve, E. 1991. "Benthic Foraminifera Reflecting Heavy Metal Pollution in Sør fjord, Western Norway." *Journal of Foraminiferal Research* 21 (1): 1-19. doi:10.2113/gsjfr.21.1.1.
- [2] Alve, E. 1995. "Benthic Foraminiferal Responses to Estuarine Pollution: A Review." *Journal of Foraminiferal Research* 25 (3): 190-203. doi:10.2113/gsjfr.25.3.190.
- [3] Dublin-Green, C. O. 1994. "Benthic Foraminifera as Pollution Indicators in the Bony Estuary, Niger Delta." *Nigerian Institute for Oceanography, Technical Paper* 95 (October): 1-21.
- [4] Moodley, L., Van Der Zwaan G. J., Herman, P. M. J., Kempers, L., and Van Breugel, P. 1997. "Differential Response of Benthic Meiofauna to Anoxia with Special Reference to Foraminifera (Protista: Sarcodina)." *Marine Ecology Progress Series* 158 (November): 151-163. doi:10.3354/Meps158151.
- [5] Le Cadre, V., and Debenay, J. P. 2006. "Morphological and Cytological Responses of Ammonia (Foraminifera) to Copper Contamination: Implication for the Use of Foraminifera as Bioindicators of Pollution." *Environmental Pollution* 143 (2): 304-17. doi:10.1016/j.envpol.2005.11.033.
- [6] Frontalini, F., and Coccioni, R. C. 2012. "The Response of Benthic Foraminiferal Assemblages to Copper Exposure: A Pilot Mesocosm Investigation." *Journal of Environmental Protection* 3 (4): 342-352. doi:10.4236/jep.2012.34044.
- [7] Ghosh, A. 2012. "Estuarine Foraminifera from the Gulf of Cambai." *Journal of the Geological Society of India* 80 (1): 65-74. doi:10.1007/s12594-012-0119-7.
- [8] Fretwell, S. D. 1972. *Populations in a Seasonal Environment*. New Jersey: Princeton University Press.
- [9] Wilson, B., and Horton, B. P. 2012. "Determining Carrying Capacity from Foraminiferal Time Series." *Journal of Micropalaeontology* 31 (November): 111-119. doi:10.1144/0262-821X11-009.

- [10] Buzas, M. A., Smith, R. K., and Beem, K. A. 1977. "Ecology and Systematics of Foraminifera in Two *Thalassia* Habitats, Jamaica, West Indies." *Smithsonian Contributions to Paleobiology* 31: 1-148.
- [11] Wilson, B. 2010. "Effects of Hurricanes on Guilds of Nearshore Epiphytal Foraminifera, Nevis, West Indies." *Journal of Foraminiferal Research* 40 (4): 327-343. doi:10.2113/gsjfr.40.4.327.
- [12] Alve, E., and Murray, J. W. 2001. "Temporal Variability in Vertical Distributions of Live (Stained) Intertidal Foraminifera, Southern England." *Journal of Foraminiferal Research* 31 (1): 12-24. doi:10.2113/0310012.
- [13] Swallow, J. E. 2000. "Intra-Annual Variability and Patchiness in Living Assemblages of Salt-Marsh Foraminifera from Mill Rytte Creek, Chichester Harbour, England." *The Micropalaeontological Society* 19 (November): 9-22. doi:10.1144/jm.19.1.9.
- [14] Murray, J. W., and Alve, E. 2000. "Major Aspects of Foraminiferal Variability (Standing Crop and Biomass) on a Monthly Scale in an Intertidal Zone." *Journal of Foraminiferal Research* 30 (3): 177-191. doi:10.2113/0300177.
- [15] Scott, D. B., and Medioli, F. S. 1980. "Quantitative Studies of Marsh Foraminiferal Distributions in Nova Scotia: Implications for Sea Level Studies." *Special Publication Cushman Foundation for Foraminiferal Research* 17: 1-58.
- [16] Sen Gupta, B. K. 1999. *Modern Foraminifera*. Netherlands: Springer.
- [17] Scott, D. B., and Medioli, F. S. 1980. "Living vs. Total Foraminifera Populations: Their Relative Usefulness in Palaeoecology." *Journal of Paleontology* 54 (4): 814-831.
- [18] Corredor, J. E., and Morell, J. M. 2001. "Seasonal Variation of Physical and Biogeochemical Features in Eastern Caribbean Surface Water." *Journal of Geophysics Research* 106 (C3): 4517-4525. doi:10.1029/2000JC000291.
- [19] Hu, C., Montgomery, E. T., Schmitt, R. W., and Muller-Karger, F. E. 2004. "The Dispersal of the Amazon and Orinoco River Water in the Tropical Atlantic and Caribbean Sea: Observations from Space and S-PALACE Floats." *Deep-Sea Research Part II: Tropical Studies in Oceanography* 51 (10-11): 1151-1171. doi:10.1016/j.dsr2.2004.04.001.
- [20] Wilson, B., Miller, K., Thomas, A., Cooke, N., and Ramsingh, R. 2008. "Foraminifera in the Mangal at Caroni Swamp, Trinidad: Diversity, Population Structure and Relation to Sea Level." *Journal of Foraminiferal Research* 38 (2): 127-136. doi:10.2113/gsjfr.38.2.127.
- [21] Horton, B. P., and Murray, J. W. 2006. "Patterns in Cumulative Increase in Live and Dead Species from Foraminiferal Time Series of Cowpen Marsh, Tees Estuary, UK: Implications for Sea-Level Studies." *Marine Micropaleontology* 58 (4): 287-315.
- [22] Horton, B. P., and Edwards, J. L. 2003. "Seasonal Distributions of Foraminifera and Their Implications for Sea-Level Studies, Cowpen Marsh, U.K." *Society for Sedimentary Geology (SEPM)* 75 (4): 21-30. doi:10.1016/S0377-8398(99)00003-1.
- [23] Walton, W. R. 1952. "Techniques for Recognition of Living Foraminifera." *Contributions from the Cushman Foundation for Foraminiferal Research* 3 (November): 56-60.
- [24] Gehrels, W. R. 1994. "Determining Relative Sea-Level Change from Salt-Marsh Foraminifera and Plant Zones on the Coast of Maine, USA." *Journal of Coastal Research* 10 (4): 990-1009.
- [25] Todd, R., and Bronnimann, P. 1957. "Recent Foraminifera and Thecamoebina from the Eastern Gulf of Paria." *Cushman Foundation for Foraminiferal Research* 3: 1-43.
- [26] Murray, J. W. 1971. *An Atlas of Recent Foraminiferids*. London: Heineman Educational Books.
- [27] Murray, J. W. 1979. *British Nearshore Foraminiferids*. London: Academic Press.
- [28] De Rijk. 1995. "Salinity Control on the Distribution of Salt Marsh Foraminifera (Great Marshes, Massachusetts)." *Journal of Foraminiferal Research* 25 (2): 156-166.
- [29] Buzas, M. A., Hayek, L. C., Reed, S. A., and Jett, J. A. 2002. "Foraminiferal Densities over Five Years in the Indian River, Lagoon, Florida: A Model of Pulsating Patches." *Journal of Foraminiferal Research* 32 (1): 68-92. doi:10.2113/0320068.
- [30] Hammer, Ø., and Harper, D. 2005. *Paleontological Data Analysis*. Oxford: Blackwell Publishing.
- [31] Hammer, Ø., Harper, D. A. T., and Ryan, P. D. 2001. "PAST: Paleontological Statistics Software Package for Education and Data Analysis." *Palaeontologia Electronica*. Accessed June 10, 2013. http://palaeo-electronica.org/2001_1/past/issue1_01.htm.
- [32] Tedesco, K. A., and Thunell, R. C. 2003. "Seasonal and Interannual Variations in Planktonic Foraminiferal Flux and Assemblage Composition in the Cariaco Basin, Venezuela." *Journal of Foraminiferal Research* 33 (3): 192-210. doi:10.2113/33.3.192.
- [33] Debenay, J. P., Guiral, D., and Parra, M. 2004. "Behaviour of Taphonomic Loss in Foraminiferal Assemblages of Mangrove Swamps of French Guiana." *Marine Geology* 208 (2): 295-314. doi:10.1016/j.margeo.2004.04.013.
- [34] Myers, E. H. 1942. "Section of Biology: Biology of the

- Foraminifera and Their Significance in Paleoecology.” *Transactions of the New York Academy of Sciences* 4 (6): 191-194. doi:10.1111/j.2164-0947.1942.tb00844.x.
- [35] Myers, E. H. 1943. “Life Activities of Foraminifera in Relation to Marine Ecology.” *Proceedings of the American Philosophical Society* 86 (July): 4389-458.
- [36] Murray, J. W. 1983. “Population Dynamics of Benthic Foraminifera: Results from the Exe Estuary, England.” *Journal of Foraminiferal Research* 13 (1): 1-12. doi:10.2113/gsjfr.13.1.1.
- [37] Erskian, M. G., and Lipps, J. H. 1987. “Population Dynamics of the Foraminiferan *Glabratella Ornatissima* (Cushman) in Northern California (USA).” *Journal of Foraminiferal Research* 17 (3): 240-256. doi:10.2113/gsjfr.17.3.240.
- [38] Ohga, T., and Kitazato, H. 1997. “Seasonal Changes in Bathyal Foraminiferal Populations in Response to the Flux of Organic Matter (Sagami Bay, Japan).” *Terra Nova* 9 (1): 33-37. doi:10.1046/j.1365-3121.1997.d01-6.x.
- [39] Lesen, A. E. 2005. “Relationship between Benthic Foraminifera and Food Resources in South San Francisco Bay, California, USA.” *Marine Ecology Progress Series* 297 (August): 131-145.
- [40] Shanks, A. L., Morgan, S. G., MacMahan, J., Reniers, A. J. H. M., Jarvis, M., Brown, J., Fujimura, A., and Griesemer, C. 2014. “Onshore Transport of Plankton by Internal Tides and Upwelling-Relaxation Events.” *Marine Ecology Progress Series* 502 (April): 39-51. doi:10.3354/meps10717.