Concentration and Biomagnification of Heavy Metals in Biota of the Coastal Marine Areas of Tanzania

Sharifa Omar Bungala, John Ferdinand Machiwa and Daniel Abel Shilla

Department of Aquatic Sciences and Fisheries, University of Dar es Salaam, Dar es Salaam 35064, Tanzania

Abstract: The concentrations of heavy metals (As, Hg, Cr, Pb and Zn) were measured in the macroalgae, macrobenthos and fish from the Tanzanian coastal marine environment in order to ascertain the biomagnification using stable isotopes of C and N. Macroalgae samples from the central marine areas of the Tanzanian coast had higher mean concentrations of Hg ($0.17 \pm 0.01 \mu\text{g/g}$) and Cr ($23.7 \pm 4.15 \mu\text{g/g}$) compared to other locations. Higher concentration of Hg ($0.06 \pm 0.02 \mu\text{g/g}$) was detected in the *Ulva fasciata* close to the Msimbazi Creek in Dar es Salaam, whereas the highest concentration of Cr ($45.5 \pm 6.83 \mu\text{g/g}$) was found in *Ulva petrusa* near Dar es Salaam port. The crab *Portunus pelagicus* collected from Pangani river estuary contained $411.5 \pm 13.04 \mu\text{g/g}$ of Zn. The other metals were uniformly distributed in macrobenthos from the entire coast. Mercury and lead in the biota were found to biomagnify along the *Arius dussumieri* and *Lethrinus lentjan* food chains as suggested by the significant positive relationships between log-pollutant concentrations in fish muscle tissues vs. $\delta^{15}\text{N}$ signatures. Zinc in muscle tissues was found to be transferred along the food webs although no biomagnification was observed. Arsenic and chromium were found to decrease with the rise of the trophic position. Metal concentrations in macroalgae, macrobenthos and fish were compared with quality guidelines values by FAO (Food and Agricultural Organization) in 1983 and they all were below permissible limits for human consumption.

Key words: Heavy metals, food chains, bioconcentration, biomagnification, stable isotope ratios, $\delta^{13}\text{C}$, $\delta^{15}\text{N}$, TL (Trophic Level).

1. Introduction

The term bioaccumulation refers to the net accumulation of a chemical by an aquatic organism as a result of uptake from all environmental sources (water, food and sediment). Bioaccumulation can be viewed as a result of competing rates of chemical uptake and elimination (chemical loss) by aquatic organisms. When the rates of chemical uptake and elimination achieve balance, the distribution of the chemical between the organism and its source(s) is said to be at steady-state [1]. Under steady-state conditions, a BAF (Bioaccumulation Factor) is defined as the ratio of the concentration of a chemical in the tissue of an aquatic organism to its concentration in water, in situations where both the organism and its food are exposed [1]. Aquatic organisms accumulate and retain certain chemicals

when exposed to these chemicals through water, their diet and sediments [2].

Biomagnification therefore is a process where chemical substances are transferred from food to an organism resulting in higher concentrations compared with the source [3-5]. Biomagnification can also be defined as the increase in concentration of chemicals between TLs (Trophic Levels). If the biomagnification factor (concentration in predator/concentration in prey) is greater than one, then the element is biomagnified in the body of a predator.

At the first TL, only direct accumulation of heavy metals from the water occurs. In considering phytoplankton, the degree of pollutant association with the cell is generally directly related to the extracellular concentration. Dead cells concentrate metals comparable to living ones indicating that the initial association of a metal with the cells is governed by adsorption [6]. Bioconcentration is the uptake of contaminant from water by aquatic organisms where water is the only pathway and is measured as a BCF

Corresponding author: Sharifa Omar Bungala, Ph.D. student, main research fields: investigation of heavy metals, organochlorines and radionuclides in aquatic food chains.

(Bioconcentration Factor). The BCF is most useful for low TL organisms such as phytoplankton and macroalgae, which are in rapid equilibrium with chemicals with low $\log K_{ow}$ (up to 5) in water [7]. BCFs do not account for biomagnification of chemicals in the food webs [8]. Hence for larger, higher TL organisms, BCF has been shown to underestimate bioaccumulation potential for hydrophobic compounds that have $\log K_{ow}$ values greater than or equal to 4.5 [8].

The role of food chains in carrying contaminants to top predator fish which is a possible source for human dietary exposure, has received attention for several decades [9-15]. The trophic transfer (trophodynamics) of elements along a food chain can result in an increase (biomagnification), a decrease (biodilution) or even may cause no change in elemental concentration along that particular food chain [16]. The trophodynamics of mercury (Hg) has been well studied and shows consistent biomagnification trends [12-15]. Other elements such as iron have been reported as not biomagnifying in food webs [9]. For other elements such as rubidium, selenium, cadmium, copper, zinc and silver, there is a growing number of studies demonstrating their biomagnification depending on site and metal-species specific factors [17-19]. Due to the deleterious effects of metals to aquatic ecosystems, it is necessary to monitor their levels in key species, because this will give an indication of the temporal and spatial extent of the accumulation and biomagnification as well as an assessment of the potential impact to human health [17].

Stable isotopes analysis has been used to study the ecological significance of materials from different sources in permanent lakes [5, 20], streams and rivers [10, 11], estuaries and bays [21, 22] and seas [23, 24]. This technique has also been used to describe trophic relationships between animals in a variety of terrestrial [25], fresh water [26] and marine [27, 28] environments. However, the great majority of these

studies have been conducted in temperate systems, and information on the sources of energy and food web structure in tropical aquatic systems in general and in estuaries of Tanzania in particular is still lacking. TP (Trophic Position) can be related with contaminant concentrations in order to trace the pathways of contaminant biomagnification by top predators in aquatic and terrestrial food webs [14, 29, 30]. Net bioaccumulation of an element occurs when its trophic transfer is greater than that of carbon and the concentration thus increases with increasing TL [31, 32].

In this study, accumulation of heavy metal As, Hg, Cr, Pb and Zn in macroalgae, macrobenthos and fish were analysed to determine biomagnification in the food chains of the marine catfish *A. dussumieri* and the snapper *L. lentjan*. The study sought to ascertain the safety of sea food from the coastal marine areas of Tanzania.

2. Materials and Methods

2.1 Study Area

This study was executed in the coastal waters of Tanzania in the western Indian Ocean. The coastal area of Tanzania (Fig. 1) encompasses a number of habitats that include coral reefs, mangroves, seagrass beds, sand banks, wetlands and beaches which support various resources both living and nonliving, and provides life support to coastal communities where such activities as fisheries and related activities play an important role in the social and economic development of local communities. The coastal areas (Dar es Salaam, Tanga, Zanzibar and Mtwara) are faced with expanding populations and emerging land-based activities such as agriculture, mineral exploitation and industrial activities which exert pressures on coastal waters, thus negatively affecting water quality. It receives water from both rivers as well as large amounts of domestic and industrial effluents. The studied area are also characterised by local harbour activities while at Dar es Salaam port

Fig. 1 Map of the marine coast of Tanzania showing the eleven sampling stations.

area, Mtwara port area and Tanga port area, there are harbour and ship fumigation activities and also loading and offloading of different types of cargo.

2.2 Establishment of the Sampling Stations

Eight sampling stations were located in the estuarine of rivers and three in marine areas near ports. Water, sediments and biota were collected from estuaries of these rivers: Pangani, Wami, Ruwusu, Mzingi, Msimbazi, Rufiji, Matandu and Ruvuma (Fig.

1). The port areas from which samples were collected were Dar es Salaam, Tanga and Mtwara (Fig. 1). The positions of sampling stations were determined using a Geographical Positioning System (Garmin GPS 72), as shown in Table 1. Three sites were established which are NCMA (Northern Coastal Marine Areas) (stations 1, 2, 3 and 4), CCMA (Central Coastal Marine Areas) (stations 5, 6 and 7) and SCMA (Southern Coastal Marine Areas) (stations 8, 9, 10 and 11).

Table 1 Sampling locations in estuaries and port areas along the coastal marine areas of Tanzania.

Site	Location	Station	Coordinates
NCMA's	TPA (Tanga Port Area)	1	5°1.43' S, 39°5.16' E
	PRE (Pangani River Estuary)	2	5°25.46' S, 38°58.01' E
	WRE (Wami River Estuary)	3	6°7.17' S, 38°48.53' E
	RRE (Ruvu River Estuary)	4	6°24.03' S, 38°52.02' E
CCMA's	MRE (Msimbazi River Estuary)	5	6°47.50' S, 39°16.54' E
	DPA (Dar es Salaam Port Area)	6	6°51.32' S, 39°17.49' E
	MZC (Mzinga Creek)	7	6°52.19' S, 39°17.21' E
	RRE (Rufiji River Estuary)	8	7°44.32' S, 39°19.11' E
SCMA's	MAE (Matandu River Estuary)	9	8°43.26' S, 39°21.41' E
	MPA (Mtwara Port Area),	10	10°16.18' S, 40°11.13' E
	RVE (Ruvuma River Estuary)	11	10°29.35' S, 40°24.58' E

2.3 Collection of Biota Samples

Biota samples were collected from eleven sampling stations along the coastal marine area of Tanzania (Fig. 1). At each station, different species of macroalgae and seagrass were collected (hand-picked) for heavy metals analysis.

Prawns and shrimps were captured during low tide with a 12-mm mesh seine net, while gastropods and crabs were also hand-picked. These macrobenthos collected from the subtidal or intertidal areas were transported to the laboratory where they were identified using procedure of Richmond, M. D. [33]. The molluscs (gastropods and bivalves) and crustaceans (crabs and shrimps) from *A. dussumieri* and *L. lentjan* food chains were dissected from their exoskeleton or shells (using dissecting kits) prior to freeze-drying. Macrobenthos tissues were freeze dried for 3 days at -60 °C and grounded into powder using a mortar and pestle.

Samples of the catfish, the emperor and some other fish were purchased from fishers in the sampling sites. Most of the fish were caught by using hook and line and some by gill netting in the estuaries and port areas. Small fish were collected along with macrobenthos using a 12-mm mesh seine net. The collected fish were counted, weighed and identified using procedure of Bianchi, G. [34] and Richmond, M. D. [33]. Total length of each fish was measured to the nearest centimeter and their wet weight established. Muscle

tissue was taken from each fish for heavy metals analysis.

For stable isotopes analysis, the tissues were freeze-dried to constant weight for three days at -60 °C and homogenized by grinding into powder using agate mortar and pestle. Using the catfish *A. dussumieri* and the emperor *L. lentjan* as examples of higher trophic consumer, all organisms that were found to be part of their food web (as deduced from the gut content analysis results) were selected for isotopic analyses from all sampling sites. The same organisms were analysed for heavy metal accumulation.

2.4 Sample Digestion and Analysis of Heavy Metals (Laboratory Analyses)

The homogenized powder of the freeze dried biota was accurately weighed (0.5 g) and digested with 5 mL of concentrated HNO₃ for 1 hour at room temperature in a Teflon beaker (covered with watch glass). The samples were then heated at 80 °C on hot plate for 4 hours. The solution was then allowed to evaporate to 2.5 mL and cooled and diluted to 50 mL with Mill-Q water [35]. The analysis of As, Hg, Cd, Cr and Zn in the samples was performed on an ICP-OES (Inductively Coupled Plasma—Optical Emission Spectroscopy), Horiba Jobin Yvon, Ultima 2 at the SEAMIC (Southern and Eastern African Mineral Centre), Dar es Salaam. The instrument was calibrated using relevant standards (Merck, Germany).

Analytical blanks were run in the similar way as that of the samples and the concentrations were determined using the standard solutions prepared in the same acid matrix. The concentration of metals in biota is expressed in dry weight basis ($\mu\text{g/g d.w.}$).

2.5 Quality Control

The accuracy of the method was checked by analysing certified reference material DORM-2 (dogfish muscle) from the National Research Council of Canada. The accuracy of the determination of heavy metals in the DORM-2 was good; it ranged from 95% to 99% (Table 2).

2.6 Statistical Analysis

Statistical analyses were performed using SPSS 16.0 for Windows, Microsoft excel and PRIMER 6. Results of the heavy metal analyses in biota were subjected to a one-way ANOVA (Analysis of Variance) to test for significant differences ($p = 0.05$) in the concentrations of the heavy metals in the tissues of the different species of biota at different sampling sites. The overall consumer isotope signatures ($\delta^{13}\text{C}$ and $\delta^{15}\text{N}$) among sites were tested by Pearson rank correlation and linear regression analysis with the concentration of heavy metals. Heavy metals concentrations were $\log_{10}$ -transformed to reduce the differences and kurtosis of the raw data prior to statistical analysis.

3. Results

3.1 Heavy Metal Concentrations in Biota Samples

3.1.1 Arsenic

The mean concentration of As in macroalgae was highest ($4.10 \pm 1.51 \mu\text{g/g}$) in samples from NCMA (stations 1, 2, 3 and 4) of Tanzania in comparison with

($2.27 \pm 0.61 \mu\text{g/g}$) in samples from CCMA (stations 5, 6 and 7) of Tanzania. The lowest concentration ($0.30 \pm 0.01 \mu\text{g/g}$) was in samples from the SCMA (stations 8, 9, 10 and 11) of Tanzania (Fig. 2). There was significant difference (ANOVA $p < 0.05$, $F = 1.023$) between As concentration in macroalgae from the three coastal locations (Table 3). The maximum concentration of As ($8.30 \pm 1.91 \mu\text{g/g}$) was measured in *S. asperifolium* from Ruvu river estuary (station 4), while the lowest concentration ($0.10 \pm 0.01 \mu\text{g/g}$) was reported in the *Chaetomorpha sp.* also from Ruvuma river estuary (station 11).

The results show that, bivalves from the CCMA had relatively higher concentrations of As ($3.17 \pm 1.22 \mu\text{g/g}$) as compared to $2.88 \pm 1.02 \mu\text{g/g}$ and $1.87 \pm 0.14 \mu\text{g/g}$ measured from SCMA and NCMA respectively (Fig. 3). There was significant difference (ANOVA $p < 0.05$, $F = 5.935$) in As contents in molluscan bivalves collected from the three coastal marine areas (Table 3). Higher concentration of As ($4.00 \pm 0.02 \mu\text{g/g}$) was found in *A. arcuatula* from Dar es Salaam port area and *Crassostrea sp.* from Ruvuma river estuary, while the lowest concentration ($1.00 \pm 0.12 \mu\text{g/g}$) was measured in *Crassostrea sp.* from Tanga port area (Fig. 3). Significantly (ANOVA, $p > 0.05$, $F = 0.457$) higher concentrations of As ($2.25 \pm 0.18 \mu\text{g/g}$) were measured in crustaceans from the NCMA while levels found in samples from CCMA and SCMA were $2.08 \pm 0.36 \mu\text{g/g}$ and $1.94 \pm 0.19 \mu\text{g/g}$ respectively (Fig. 4). The highest concentration of As ($4.00 \pm 1.11 \mu\text{g/g}$) was measured in the shrimp *P. indicus* from Msimbazi river estuary, while the lowest concentration ($0.50 \pm 0.02 \mu\text{g/g}$) was found in the crab *P. pelagicus* from Matandu river estuary (Fig. 4).

Levels of As recorded in fish species from the CCMA was relatively higher ($3.40 \pm 1.03 \mu\text{g/g}$)

Table 2 Mean ($\pm$ SD) concentration of metals ($\mu\text{g/g}$) in the certified reference material DORM-2 (SD: Standard Deviation).

	As	Hg	Cr	Pb	Zn
Certified	18 ± 1.7	4.64 ± 0.26	34.7 ± 5.5	0.065 ± 0.007	25.6 ± 2.3
Observed	17.5 ± 3.2	4.43 ± 0.27	34.4 ± 3.2	0.062 ± 0.009	25.3 ± 1.9
% recovery	97.2	95.4	99.1	95.5	98.8

Fig. 2 Concentration (mean $\pm$ SD) of As in biota samples (A = macroalgae; B = bivalves; C = crustaceans; D = fish) from the NCMA, the CCMA and the SCMA.

compared to 2.08 ± 0.97 $\mu\text{g/g}$ and 0.62 ± 0.14 $\mu\text{g/g}$ in fish species from SCMA and NCMA respectively (Fig. 5). As content (7.80 ± 2.03 $\mu\text{g/g}$) found in the zoobenthivorous fish *P. quadrilineatus* from Mtwara

port area (station 10) was significantly higher than concentrations measured in the omnivorous fish *A. dussumieri* (0.10 ± 0.04 $\mu\text{g/g}$) from Matandu river estuary (ANOVA $p < 0.05$, $F = 0.583$) (Table 3). As concentrations in fish were compared with quality guidelines values FAO (Food and Agricultural Organization) [36] and they all were below permissible limits of 1.4 $\mu\text{g/g}$ w.w. (Table 4).

3.1.2 Mercury

The mean concentration of mercury in macroalgae samples was almost similar (0.03 ± 0.01 $\mu\text{g/g}$) in samples from SCMA, NCMA and CCMA. There was significant difference (ANOVA $p < 0.05$, $F = 1.286$) in Hg concentration in macroalgae from the three coastal marine areas of Tanzania (Table 3). The maximum concentration of Hg (0.06 ± 0.03 $\mu\text{g/g}$) was found in the *U. fasciata* from Msimbazi river estuary, while the lowest concentration of the metal ($0.01 \pm$

Table 3 ANOVA, heavy metal concentrations in macroalgae, macrobenthos, crustaceans and fish samples.

	Sum of squares	Df *	Mean square	F	Sig.
Macroalgae					
As	38.031	7	5.433	1.023	0.544
Hg	0.002	7	0	1.286	0.457
Cr	1,774.511	7	253.502	5.778	0.089
Pb	10.259	7	1.466	0.402	0.855
Zn	13,789.41	7	1,969.916	0.28	0.925
Bivalves					
As	10.386	7	1.484	5.935	0.086
Hg	0.001	7	0	1.223	0.476
Cr	354.784	7	50.683	2.153	0.284
Pb	1,787.67	7	255.381	3.201	0.184
Zn	1,054,670	7	150,667.1	4.631	0.118
Crustaceans					
As	5.818	10	0.582	0.457	0.886
Hg	0	10	0	0.189	0.993
Cr	41.25	10	4.125	9.811	0
Pb	2.704	10	0.27	1.565	0.237
Zn	119,718.8	10	11,971.88	2.428	0.081
Fish					
As	111.747	26	4.298	0.583	0.843
Hg	0.018	26	0.001	3.844	0.05
Cr	594.492	26	22.865	0.323	0.98
Pb	90.464	26	3.479	11.134	0.003
Zn	420,978.8	26	16,191.49	149.202	0

*Df: degree of freedom.

Fig. 3 Concentration (mean $\pm$ SD) of Hg in A = macroalgae; B = bivalves; C = crustaceans; and D = fish from the NCMA, CCMA and SCMA.

Table 4 The average concentrations of trace metals in fish as recommended by the FAO [36].

Specie	Unit	Pb	Zn	Cr	Hg	As
Fish	$\mu\text{g/g w.w.}$	2	50	1	0.5	1.4

0.00 $\mu\text{g/g}$) was found in the *U. petrusa* from the Dar es Salaam port area (Fig. 2). Bivalves from the CCMA had relatively higher concentrations of Hg ($0.04 \pm 0.02 \mu\text{g/g}$) compared to the concentration ($0.04 \pm 0.01 \mu\text{g/g}$) in NCMA and concentration ($0.03 \pm 0.01 \mu\text{g/g}$) in SCMA. There was significant difference (ANOVA $p < 0.05$, $F = 1.223$) in Hg contents in bivalves collected from the three main coastal locations (Table 3). The highest concentration of Hg ($0.06 \pm 0.03 \mu\text{g/g}$) was found in *A. arcuatula* from Mzingu Creek, while the lowest concentration ($0.02 \pm 0.01 \mu\text{g/g}$) was found in *Crassostrea sp.* from Ruvuma river estuary (Fig. 3).

The results show that crustaceans from the CCMA had almost similar concentrations of Hg ($0.04 \pm 0.02 \mu\text{g/g}$) with NCMA. Lower concentration ($0.03 \pm 0.01 \mu\text{g/g}$) was found in SCMA. There was significant difference (ANOVA $p < 0.05$, $F = 0.189$) in Hg contents in crustaceans collected from the three coastal locations (Table 3). Higher concentration of Hg ($0.05 \pm 0.03 \mu\text{g/g}$) was found in the crab *P. pelagicus* from Dar es Salaam port and Tanga port areas and in the crab *S. serrata* from the estuaries of

Wami and Ruvu rivers. Lower concentration of Hg ($0.02 \pm 0.01 \mu\text{g/g}$) was found in the shrimp *P. indicus* from stations 1, 3, 7, 10 and 11 (Fig. 4). Like crustaceans, fish species from CCMA had almost similar concentrations of Hg ($0.05 \pm 0.03 \mu\text{g/g}$) as those from SCMA ($0.05 \pm 0.02 \mu\text{g/g}$). The lower concentration ($0.04 \pm 0.01 \mu\text{g/g}$) was reported in NCMA (Fig. 4). There was significant difference (ANOVA $p < 0.05$, $F = 3.844$) in Hg contents in the fish species collected from the three coastal locations (Table 3). Higher concentration of Hg ($0.08 \pm 0.03 \mu\text{g/g}$) was found in the carnivorous fish *L. lentjan* from Msimbazi river estuary and also in the tissues of an omnivorous fish *A. dussumieri* from Matandu river estuary (Fig. 5). The lowest concentration ($0.01 \pm 0.00 \mu\text{g/g}$) was found in the herbivorous fish *S. ghobban* from Tanga port area, planktivorous fish *H. kelee* from Ruvu river estuary, planktivorous fish *S. albella* from Matandu river estuary and in the detritivorous fish *V. seveli* from Ruvuma river estuary (Fig. 4). Hg concentrations in fish were compared with quality guidelines values FAO [36] and they all were below permissible limits of $0.5 \mu\text{g/g w.w.}$ (Table 4).

3.1.3 Chromium

The highest concentration of chromium was found in the macroalgae samples ($23.67 \pm 8.19 \mu\text{g/g}$) from the CCMA of Tanzania. An intermediate concentration ($12.00 \pm 4.56 \mu\text{g/g}$) was found in the NCMA of Tanzania, while lowest concentration ($3.75 \pm 1.81 \mu\text{g/g}$) was found in the samples from the SCMA of Tanzania (Fig. 3). These concentrations were statistically different across the three coastal locations (ANOVA, $p < 0.05$, $F = 5.778$) (Table 3). The maximum concentration of Cr ($45.50 \pm 6.38 \mu\text{g/g}$) was found in the *U. petrusa* from Dar es Salaam port area, while the lowest concentration ($2.00 \pm 0.03 \mu\text{g/g}$) was found in the *Chaetomorpha sp.* from Ruvuma river estuary (Fig. 2). The bivalves from the NCMA had relatively higher concentration of Cr ($11.13 \pm 5.25 \mu\text{g/g}$) compared to $7.40 \pm 2.14 \mu\text{g/g}$ and $5.00 \pm 1.01 \mu\text{g/g}$ recorded in from SCMA and CCMA

respectively (Fig. 3). Although there was significant difference in mean Cr contents in molluscan bivalves collected from the three coastal locations (ANOVA, $p < 0.05$, $F = 2.153$), within site comparison revealed that higher concentration ($21.00 \pm 9.19 \mu\text{g/g}$) was found in *S. cuculata* from Mtwara port area while lower concentrations ($2.00 \pm 0.42 \mu\text{g/g}$) were found in the tissues of the *A. arcuatula* and *Crassostrea sp.* from Mzinga Creek and Ruvuma river estuaries respectively (Fig. 3 and Table 3). Crustaceans from the SCMA had relatively higher concentrations of Cr ($3.10 \pm 0.88 \mu\text{g/g}$) compared to $3.00 \pm 1.14 \mu\text{g/g}$ in samples from CCMA and $2.25 \pm 1.33 \mu\text{g/g}$ in samples from NCMA. There was no significant difference (ANOVA $p > 0.05$) in Cr contents in crustaceans collected from the three coastal marine areas (Table 3). The highest concentration of Cr ($8.00 \pm 3.17 \mu\text{g/g}$) was found in the shrimp *P. indicus* from Mtwara port area, while the lowest concentration ($1.00 \pm 0.02 \mu\text{g/g}$) was found in the crab *S. serrata* from Rufiji river estuary (Fig. 4). Results further elucidate that fish species from the NCMA accumulated relatively higher Cr ($5.50 \pm 0.19 \mu\text{g/g}$) compared to values recorded in fish species from CCMA ($4.80 \pm 0.61 \mu\text{g/g}$) and SCMA ($2.30 \pm 0.13 \mu\text{g/g}$). There was significant difference in Cr contents in the fish species collected from the three coastal locations (ANOVA $p < 0.05$, $F = 0.323$) (Table 3). The highest concentration of Cr ($14.50 \pm 5.13 \mu\text{g/g}$) was found in the carnivorous fish *L. lentjan* collected from Mzinga Creek, while the lowest concentration ($1.00 \pm 0.02 \mu\text{g/g}$) was found in the detritivorous fish *L. macrolepis* from Rufiji river estuary and herbivorous fish *S. sutor* from Dar es Salaam port area (Fig. 5). Cr concentrations in fish were compared with quality guidelines values FAO [36] and they all were below permissible limits of $1.0 \mu\text{g/g w.w.}$ (Table 4).

3.1.4 Lead

The mean concentration of lead in macroalgae samples was highest ($2.75 \pm 0.52 \mu\text{g/g}$) in samples from the NCMA of Tanzania, followed by the

Fig. 4 Concentration (mean $\pm$ SD) of Cr in A = macroalgae; B = bivalves; C = crustaceans; and D = fish from NCMA, CCMA and SCMA.

Fig. 5 Concentration (mean $\pm$ SD) of Pb in A = macroalgae; B = bivalves; C = crustaceans; and D = fish from NCMA, CCMA and SCMA.

CCMA ($1.37 \pm 0.18 \mu\text{g/g}$) and $1.15 \pm 0.26 \mu\text{g/g}$ in samples from the SCMA of Tanzania (Fig. 2). Significant difference existed in mean Pb concentrations of macroalgae samples collected from the three coastal locations (ANOVA $p < 0.05$, $F = 0.402$), (Table 3). Maximum concentration of Pb ($5.50 \pm 1.08 \mu\text{g/g}$) was found in *S. asperifolium* from Ruvu river estuary, while the minimum concentration of the metal ($0.10 \pm 0.02 \mu\text{g/g}$) was measured in the *U. petrusa* from Dar es Salaam port area. The results also show that bivalves from the CCMA had relatively higher concentrations of Pb ($2.83 \pm 0.86 \mu\text{g/g}$) compared to bivalves from NCMA ($2.05 \pm 0.93 \mu\text{g/g}$) and SCMA ($1.40 \pm 0.65 \mu\text{g/g}$). Significant difference

was found in Pb contents in molluscan bivalves across the three coastal locations (ANOVA $p < 0.05$, $F = 3.201$). Highest concentration of Pb ($5.00 \pm 1.17 \mu\text{g/g}$) was found in *Crassostrea sp.* from Pangani river estuary and lower values ($0.50 \pm 0.02 \mu\text{g/g}$) was measured in *Crassostrea sp.* from Tanga port area (Fig. 5). Crustaceans from the SCMAS had relatively higher concentrations of Pb ($1.24 \pm 0.52 \mu\text{g/g}$) compared to crustaceans collected from NCMAAs ($1.03 \pm 0.46 \mu\text{g/g}$) and CCMAAs ($0.92 \pm 0.31 \mu\text{g/g}$) (Fig. 3). Statistical results revealed significant difference in Pb content of crustacean tissues across the three sampled coastal marine areas (ANOVA $p < 0.05$, $F = 1.565$). Higher concentration of Pb ($1.80 \pm 0.39 \mu\text{g/g}$) was measured in the crab *P. pelagicus* from Ruvuma river estuary while the shrimp *P. indicus* from Wami river estuary, Ruvuma river estuary and Mtwara port area; and the crab *S. serrata* from Wami river estuary and *P. pelagicus* from Mzinga Creek recorded lower concentration ($0.50 \pm 0.02 \mu\text{g/g}$) of lead (Fig. 4). Results further reveal that fish species from the CCMAAs had relatively higher average concentrations of Pb ($2.60 \pm 1.32 \mu\text{g/g}$) compared to $2.33 \pm 0.98 \mu\text{g/g}$ from SCMAAs and $2.29 \pm 1.05 \mu\text{g/g}$ from NCMAAs. These results were statistically significant different across the three coastal marine locations (ANOVA $p < 0.05$, $F = 11.134$). The highest concentration of Pb ($8.00 \pm 3.07 \mu\text{g/g}$) was measured in the carnivorous *L. lentjan* from Dar es Salaam port area, while the lowest concentration ($0.50 \pm 0.12 \mu\text{g/g}$) was found in the herbivorous *S. ghobban* from Tanga port area, the planktivorous *S. albella* from the estuaries of Wami and Ruvu rivers, the detritivorous *L. macrolepis* and *V. seheli* from the estuaries of Rufiji and Ruvuma rivers respectively (Fig. 5). Pb concentrations in fish were compared with quality guidelines values FAO [36] and they all were below permissible limits of $2.0 \mu\text{g/g}$ w.w. (Table 4).

3.1.5 Zinc

The average concentration of zinc in macroalgae samples was highest ($111.50 \pm 5.17 \mu\text{g/g}$) in samples

from NCMAAs followed by SCMAAs ($60.63 \pm 3.53 \mu\text{g/g}$) and $51.00 \pm 1.26 \mu\text{g/g}$ in samples from CCMAAs. These results illustrate similar distribution of this metal across the three coastal marine areas (ANOVA $p < 0.05$, $F = 0.280$) (Table 3). Inter-site comparison show that highest concentration of Zn ($216.00 \pm 11.66 \mu\text{g/g}$) was found in *S. asperifolium* from Ruvu river estuary, while the lowest concentration ($23.50 \pm 8.93 \mu\text{g/g}$) was measured in the *U. fasciata* from Pangani river estuary (Fig. 2). The bivalves from the CCMAAs recorded higher concentrations of Zn ($616.50 \pm 21.14 \mu\text{g/g}$) compared to $167.30 \pm 16.85 \mu\text{g/g}$ from NCMAAs and $116.1 \pm 18.22 \mu\text{g/g}$ from SCMAAs. There was no significant difference (ANOVA $p > 0.05$, $F = 4.631$) in Zn contents in molluscan bivalves collected from the three coastal locations (Table 3). The highest concentration of Zn ($651.00 \pm 6.51 \mu\text{g/g}$) was measured in *Crassostrea sp.* from Msimbazi river estuary, while the lowest concentration ($29.50 \pm 7.93 \mu\text{g/g}$) was found in *S. cucullata* from Mtwara port area (Fig. 3).

Crustaceans from the NCMAAs had relatively higher concentrations of Zn ($165.90 \pm 33.39 \mu\text{g/g}$) compared to the concentration ($65.10 \pm 15.16 \mu\text{g/g}$) in CCMAAs and the lowest concentration ($52.00 \pm 22.16 \mu\text{g/g}$) in samples from SCMAAs. There was significant difference (ANOVA $p < 0.05$, $F = 2.428$) in Zn content in crustaceans collected from the three coastal locations (Table 2). The highest concentration of Zn ($411.50 \pm 13.04 \mu\text{g/g}$) was found in the crab *P. pelagicus* from Pangani river estuary, while the lowest concentration ($28.00 \pm 4.34 \mu\text{g/g}$) was measured in the shrimp *P. indicus* from Mtwara port area (Fig. 4). The results show that fish species from the SCMAAs had higher mean concentration of Zn ($101.10 \pm 21.16 \mu\text{g/g}$) compared to NCMAAs ($84.00 \pm 15.03 \mu\text{g/g}$) and CCMAAs ($61.40 \pm 19.82 \mu\text{g/g}$). There was no significant difference (ANOVA $p > 0.05$) in mean Zn contents in the fish species collected from the three coastal marine areas (Table 3). The highest concentration of Zn ($667.00 \pm 43.11 \mu\text{g/g}$) was measured in the omnivorous

Fig. 6 Concentration (mean $\pm$ SD) of Zn in A = macroalgae; B = bivalves; C = crustaceans; and D = fish from NCMA, CCMA and SCMA.

fish *A. dussumieri* from Matandu river estuary, while the lowest concentration (17.00 ± 9.48 µg/g) was found in the zoobenthivorous fish *P. quadrilineatus* from Mtwara port area (Fig. 6). Zn concentrations in fish were compared with quality guidelines values FAO [36] and they all were below permissible limits of 50.0 µg/g w.w. (Table 4).

3.2 The TPs of *A. dussumieri* and *L. lentjan*

The estimation of the positions of the cat fish *A. dussumieri* and the emperor *L. lentjan* in the estuarine and nearshore waters of Tanzania was made using the stable C and N isotope results. The TP of fishes (TP_{SIA}), was calculated following the procedure of Cabana, G. and Rasmussen, J. B. [37] using Eq. (1):

$$TP_{SIA} = \left[\left(\text{Fish } \delta^{15}N - \text{Primary consumer } \delta^{15}N \right) / 3.4 \right] + 2 \quad (1)$$

Primary consumer $\delta^{15}N$ was calculated as the mean $\delta^{15}N$ of all primary consumers (i.e., filter feeder, collector, gatherer, shredder and grazer taxa) collected for a sampling event. Trophic guilds were assigned using the literature [38]. Stomach contents analysis also enabled the determination of TP of *A. dussumieri* and *L. lentjan*. By using gut content analysis, the TP of fish (TP_{GCA}) was also calculated according to the method of Vander-Zanden, M. J., et al. [39] using Eq. (2):

$$TP_{GCA} = \sum (P_i * T_i) \quad (2)$$

Where P_i is proportion of the i^{th} food item and T_i is TP of each food item. TPs of food items were based on literature dietary data [39, 40]. According to Lopez-Fernandez, H. and Winemiller, K. O. [38], in marine systems there are: algae at the bottom of the food web (TL = 1, by definition); herbivorous zooplankton feeding on the algae (TL = 2); large zooplankton or small fishes, feeding on the herbivorous zooplankton (TL = 3); large fishes (for example, cod, tuna and groupers) whose food tends to be a mixture of low-TP and high-TP organisms (TL = 3.5-4.5).

The stable nitrogen isotopic signatures results show that *A. dussumieri* and *L. lentjan* from all sampling stations belong to the TL of the secondary consumers. The estimated TP based on the stable isotopes of carbon and nitrogen based on the procedure of Cabana, G. and Rasmussen, J. B. [37] for *A. dussumieri* was 3.5 and that of *L. Lentjan* was 3.0. Individuals of *A. dussumieri* and *L. lentjan* increased in weight linearly, as their length increased. These findings are in agreement with the TP of marine fishes as estimated using $\delta^{15}N$ values [41]. Cabana, G. and Rasmussen, J. B. [37] reported a linear relationship between the size of individuals of benthic assemblages and their stable nitrogen isotope values. The good agreement between the determination of the TLs of 3.5 and 3.0 for *A. dussumieri* (Fig. 7) and *L. lentjan* (Fig. 8) respectively by using $\delta^{15}N$ method and the traditional approach of using the gut-content analysis supports the application of an average trophic enrichment of 3.4‰ between successive TL in the estimation of TPs of consumers as suggested by Vander-Zanden, M. J. and Rasmussen, J. B. [42].

3.3 Trophic Transfer of Heavy Metals along the Food Chains

The stable isotopes of C and N ($\delta^{13}C$ and $\delta^{15}N$) for the *A. dussumieri* and *L. lentjan* and their TLs were used in the investigation of trophic transfers of heavy metals along their food chains.

Fig. 7 Relationship between $\delta^{15}\text{N}$ and log-transformed Hg concentration in fish from Tanzanian coastal marine environment.

Fig. 8 Relationship between $\delta^{15}\text{N}$ and log-transformed Pb concentration in fish from Tanzanian coastal marine environment.

Fig. 9 Relationship between $\delta^{15}\text{N}$ and log-transformed As concentration in fish from Tanzanian coastal marine environment.

Fig. 12 Relationship between $\delta^{15}\text{N}$ and log-transformed Pb concentration in fish from Tanzanian coastal marine environment.

Fig. 10 Relationship between $\delta^{15}\text{N}$ and log-transformed Hg concentration in fish from Tanzanian coastal marine environment.

Fig. 13 Relationship between $\delta^{15}\text{N}$ and log-transformed Zn concentration in fish from Tanzanian coastal marine environment.

Fig. 11 Relationship between $\delta^{15}\text{N}$ and log-transformed Cr concentration in fish from Tanzanian coastal marine environment.

In the *A. dussumieri* and *L. lentjan* food chains, log concentrations of mercury and lead in muscle and soft tissues of animal samples were found to increase significantly ($p = 0.05$ and $p = 0.01$) with increasing TLs (Figs. 9 and 10). The slopes for these elements were 0.368 for log-transformed Hg and 0.246 for log-transformed Pb. Regressions for log As and Cr were significant, with a negative slope of -0.068 and -0.005 respectively (Figs. 11 and 12). Non-significant slope was found for Zn, indicating that this element neither biomagnifies nor biodilutes through the food web (Fig. 13).

There was a linear relationship between

log-transformed Hg and Pb concentrations with $\delta^{15}\text{N}$ values for biota analysed. The slope of regression line was negative between log-transformed As and Cr concentrations with $\delta^{15}\text{N}$ values, while completely flat regression line was between log-transformed Zn concentration with $\delta^{15}\text{N}$ values for biota. The steeper slope was observed between log-transformed Hg with $\delta^{15}\text{N}$ in biota as compared to the slope for Pb.

4. Discussion

4.1 Accumulation and Biomagnification of Heavy Metals along Food Chains of *A. dussumieri* and *L. lentjan*

4.1.1 Arsenic

Bioaccumulation of As was high in macroalgae from NCMAs followed by SCMAS and the last was CCMAs. Although As was not detected in the water samples, it was found to bioaccumulate in the three species of macroalgae. The significant BCF for As in the macroalgae *S. asperifolium* in Ruvu river estuary and *U. fasciata* in Msimbazi Creek was expected because of the high concentration of the As in the sediments from the stations. Higher BCF for As in *U. fasciata* from Wami river estuary could be due to high ability of *Ulva* sp. to accumulate arsenic from the environment as compared to other species of macroalgae in other stations. This idea is complemented by discussion given by Thomson, D., et al. [43] who reported that arsenic concentration and which are important primary producer in marine food chains. Rahman, M. A., et al. [44] reported that total arsenic concentrations varied between classes of algae, and significant differences between algae classes and habitats were found for the proportion of As species.

Bioaccumulation of arsenic in bivalves was generally low. The only significant BAF was found in *S. cucullata* in Mtwara port area (BAF = 133) in SCMAS. The results show that no significant BAF was found in crustaceans from NCMAs, CCMAs and also from SCMAS. The findings are in agreement with the previous report by Barwick, M. and Maher, W. [45]

which had reported that As and arsenic-containing organic compounds did not accumulate to a great extent in aquatic organisms due to its highly toxic nature.

The results on As bioaccumulation in fish showed that the metal had relatively high BAF in the fish from CCMAs, followed by SCMAS and then NCMAs. Only five fish species were found with significant BAF showing that they accumulate As from the environment. These fish species include the herbivorous *S. sutor* from Msimbazi river estuary (BAF = 108), the zoobenthivorous *P. quadrilineatus* from Mzinga Creek (BAF = 300), the omnivorous *A. dussumieri* from Mtwara port area (BAF = 183), and Wami river estuary (BAF = 133) as well as the zooplanktivorous *S. albella* also from Wami river estuary (BAF = 109). The high BAF for these fish suggests high availability of As at stations 3, 5 and 7. Hardersen, S. and Wratten, S. D. [46] reported in their research the uptake of As by fish from water, food, sediments and suspended particulate material. Therefore, arsenic present in both water and sediments at these sampling stations was available for uptake by fish. The uptake of sediment-associated contaminants by fish may occur by respiratory and dietary routes, whereas the dermal route is usually a minimal contributor of exposure, due to the often effective barrier provided by the external epithelium [47].

The result of trophic transfer (trophodynamics) of As along *A. dussumieri* and *L. lentjan* food chain is described by negative regression which means biodilution of the metal (Fig. 8). The finding was confirmed by the elevated concentrations of arsenic in the macroalgae, macrobenthos and small fish, while larger fish had lower concentrations. Interpretation of this observation is supported by the findings of Wang, W. X. [9] who reported that concentrations of metals decreased at a higher TL in the marine food chain. This is mainly a result of an effective efflux of metals from macrobenthos and a very low assimilation of metals by marine fish. Other researchers have reported

that arsenic and/or its metabolites is a chemical that bioaccumulates in tissues of aquatic organisms but does not biomagnify, rather it biodilutes in the aquatic food chain [31, 48-50]. Yet Thomson, D., et al [43] report that, dietary exposure to arsenic from aquatic foods would not be a serious problem for humans due to its biodiminution and biotransformation to less toxic organo-As species.

4.1.2 Mercury

Results show that bioconcentration of mercury were higher in the macroalgae from CMCA's followed by SCMA's and NCMA's. Bioaccumulation of mercury was higher in the macroalgae than in all other biota analysed, with significant BCFs measured in eight stations out of eleven. Higher significant BCF values were found in *U. fasciata* from Msimbazi river estuary (1,700), in *U. petrusa* from Dar es Salaam port area (700) and in *Chaetomorpha sp.* from Rufiji river estuary (900). These results show that *Ulva sp.* and *Chaetomorpha sp.* have a higher ability to accumulate Hg from the environment as compared to other macroalgae species. The high BCFs for macroalgae were also related to the sampling stations with higher concentrations of available Hg in water and sediments. Karez, C. S., et al. [51] reported that high levels of Hg in the algae reflect the high bioavailability of metals and the capacity of the algae to accumulate metals.

The macrobenthos (bivalves and crustaceans) had generally low BAFs throughout the study area suggesting that most of the macrobenthos accumulate heavy metals from biota they consume. Low BAFs values for this group of organisms could, therefore, be due the use of water and sediments data in computing BAFs. Similar results have been reported by Lee, B. G., et al. [52] that most benthic organisms obtain a large fraction of their contaminants through ingestion and not by direct accumulation from overlying water or pore-water. Similarly, the BAFs for mercury in fish from coastal marine area of Tanzania were low and not significant. These indicative results suggest that at present, fish in the coastal marine areas of Tanzania

are not bioaccumulating Hg from the environment. Low BAFs for Hg in fish are probably a result of the low concentration of available Hg in these environments. Romeo, M., et al. [53] observed that the ability of fish to accumulate heavy metals depended on ecological needs, metabolism, and the extent of pollution in sediment, water and food items.

Results on trophic transfer of Hg along *A. dussumieri* and *L. lentjan* food chains in this study show that biomagnification of Hg was evidenced by a positive regression results (Fig. 9). This was further confirmed by elevated Hg concentrations in the big fish, while small fish, macroalgae and macrobenthos had lower concentrations. Regression relationships between Hg concentrations and $\delta^{15}\text{N}$ have been discussed in other aquatic food chains [15, 29, 30, 54, 55] demonstrating biomagnification of mercury along food chains despite low ambient concentrations of this metal in the environment. Results of this study display similar mechanism of biomagnification of metals from lower TL organisms such as plankton or algae to higher TL organisms such as *A. dussumieri* and *L. lentjan* food webs.

Higher Log.-mercury slope found in this study compared to Log.-lead slope gives evidence that Hg is the major biomagnifying element in *A. dussumieri* and *L. lentjan* food chains from Tanzania coastal marine environment. This means that Hg is biomagnified through this food webs as evidenced by $\delta^{15}\text{N}$ regardless of productivity or salinity (marine vs. freshwater) of the ecosystem. This argument is supported by the results reported by Campbell, L. M., et al. [15] who found that MeHg measured by $\delta^{15}\text{N}$ is biomagnified through diverse food webs, regardless of productivity (eutrophic vs. oligotrophic), latitude (arctic vs. tropical) or salinity (marine vs. freshwater) of the ecosystem.

4.1.3 Chromium

The results show that the bioconcentration of chromium was higher in the macroalgae from CCMA's, followed by macroalgae from NCMA's and SCMA's.

Generally, chromium was found to significantly bioaccumulate in all the analysed species of macroalgae. Significant BCF values for Cr were found in the *U. petrusa* from Dar es Salaam port area (1,820) and in *U. fasciata* from Wami river estuary (1,240). The high BCFs for macroalgae are in agreement with the high Cr concentration that was observed in the water samples from the sampling stations. *Ulva sp.* had the highest ability to accumulate Cr as compared to the other species of macroalgae. The high BCFs are also indicative of the high dissolved Cr in water samples from Wami river estuary and Dar es Salaam port area. According to Luoma, S. N. [56], unlike other bioindicators of heavy metal contamination such as filter-feeding animals, macroalgae accumulate dissolved metal ions in the waters. Therefore, only heavy metals with high solubility products are expected to have significant BCFs in macroalgae.

The bivalves from NCMA had high BAFs for chromium, followed by SCMA and CCMA. Significant BAFs for Cr in bivalves were measured only in *S. cucullata* from Wami river estuary (220), Ruvu river estuary (111) and Mtwara port area (300). Similarly, water samples from CCMA were found to contain higher particulate chromium as well as higher percentage of available Cr in the sediment as compared to NCMA and SCMA. *S. cucullata* was found to have high ability to accumulate and store chromium in their tissues as compared to the ability of *A. arcuata*. This observation is in agreement with the research data from China in which *S. cucullata* were described as the strongest metal accumulators among bivalve species [57]. The accumulation of chromium by crustaceans from the coastal marine areas of Tanzania was low with no crustacean found having significant BAF. Similarly, there was a low bioaccumulation of Cr in fish from the studied locations with no fish having significant BAF. It can therefore be suggested that crustaceans and fish do not significantly bioaccumulate chromium from the environment (water and sediment).

The biomagnification results showed that there was a decrease of concentration of Cr along *A. dussumieri* and *L. lentjan* food chains (biodilution) which was implied by negative regressions graphs (Fig. 10). The result was further confirmed by the elevated concentrations of Cr in the macroalgae, macrobenthos and small fish, while larger fish had lower concentrations. It has been reported that concentrations of metals decrease at a higher TL in the marine food chain as a result of an effective efflux of metals from macrobenthos and a very low assimilation of metals by marine fish [9].

4.1.4 Lead

Lead is the only heavy metal which has no significant BCF in macroalgae from all the three main sites despite some high Pb concentrations in sediments from the study area. The low BCFs in the macroalgae are related to the low concentration of lead in the water samples which was largely below the detection limit. It is well known that macroalgae accumulate heavy metals proportional to the surrounding medium [58] and that the accumulation of the heavy metal such as Pb in macroalgae depends on the factors like metal contents in water, absorption properties of the algae species, and to some extent the metal content of sediment [59].

Bivalves *S. cucullata* from the estuaries of Rufiji and Matandu rivers (290 and 193, respectively) in SCMA had significant BAF for lead. Significant BAF was also found in *S. cucullata* from Wami river estuary (133) and Ruvu river estuary (141) in NCMA. Other bivalves displayed insignificant values for BAF of this metal. On the other hand, sediments from the estuaries of Wami, Ruvu, Rufiji and Matandu rivers had high concentration of available Pb which possibly has an influence on the observed BAFs. Insignificant BAFs displayed by crustaceans and fish are a clear indication that these species do not substantially bioaccumulate Pb from the coastal marine areas of Tanzania.

Positive regressions result elucidates that Pb biomagnifies along *A. dussumieri* and *L. lentjan* food

chains in Tanzanian coastal marine waters (Fig. 11). Elevated Pb concentrations in the big fish species relative to small fish, macroalgae and macrobenthos displaying lower concentrations of this metal in this study further confirm this observation. Positive regression relationships between $\log_{10}$ transformed Pb concentrations and $\delta^{15}\text{N}$ have been used by different authors in delecting biomagnification of this metal in aquatic food webs [12, 18, 29, 30]. In the present study, the graph (Fig. 11) of log-transformed Pb slope was lower than that of Hg indicating that Hg potentially biomagnifies more than Pb along *A. dussumieri* and *L. lentjan* food chains.

4.1.5 Zinc

The results show high BCF for zinc in macroalgae from the coastal marine areas of Tanzania. Highly significant BCF values were observed in *S. asperifolium* from Ruvu river estuary (3,323), in *Chaetomorpha sp.* from Ruvuma river estuary (1,536), *U. fasciata* from Matandu river estuary (1,164) and in *U. fasciata* from Wami river estuary (1,088). The high BCFs for macroalgae are consistent with the observed high zinc concentration in the macroalgae. The results also suggest that the macroalgae have a high ability of bioaccumulating zinc in the tissue.

The bioaccumulation of Zn was also high in bivalves from all sites. The significant BAFs were found in *A. arcuatula* from Mzinga Creek (1,630) and *S. cuculata* from the estuaries of Ruvu, Wami and Rufiji rivers (307, 204 and 187, respectively). The high values of BAFs relates well with the concentration of Zn in the suspended particulate matter which was higher at Mzinga Creek and Rufiji river estuary. Similarly, higher concentrations of available zinc in sediments were found at the estuaries of Msimbazi, Rufiji and Ruvu rivers. The current study argues that bivalves have high ability to accumulate zinc, and that molluscan life processes do not seem to be affected by excess zinc which is frequently accumulated far in excess of the organism's immediate needs [60].

The bioaccumulation of zinc in crustaceans from the coastal marine areas of Tanzania was high. The significant BAF was observed in the *P. indicus* and *P. pelagicus* from Mtwara port area, in *P. indicus* and *P. pelagicus* from Ruvuma river estuary and in *P. indicus* and *S. serrata* from Rufiji river estuary. However, there was no crustacean with significant BAF from NCMA's, probably due to the lower values of available Zn in the sediment from NCMA's. A study by Romeo, M., et al. [61] discusses that Zn can be taken up and accumulated in tissues and bodies of many species of marine invertebrates to concentrations usually much higher than concentrations in the surrounding seawater or sediments.

Only four fish species were found to accumulate zinc in their tissues to a significant extent from the coastal marine areas of Tanzania. The significant BAF was observed in the omnivorous *A. dussumieri* from the estuaries of Matandu, Ruvu and Wami rivers (733, 364 and 146, respectively) and Mtwara port area (143). The observed significant BAF of Zn in the catfish *A. dussumieri* from all sites was probably due to the feeding habit. Previous studies [62, 63] have reported that variability of levels of metals such as Zn in different fish species is a function of feeding habits, ecological needs, metabolism or age, size and length of the fish and their habitats, which affirms the argument.

Zinc trophodynamic results are explained by positive regression slope and confirmed by evenly distribution of Zn in macroalgae, macrobenthos and fish. The results clarify that Zn concentrations did not increase or decrease with TLs along *A. dussumieri* and *L. lentjan* food chains (Fig. 12). Zinc is an essential element with many important metabolic functions which necessitates most organisms to have biochemical mechanisms to regulate the amount of Zn in their cells [63]. Campbell, L. M., et al. [15], suggested that the rate of Zn excretion would be relative to Zn ingestion, thereby ensuring a relatively constant concentration in biota throughout the food web.

5. Conclusion

This study revealed evidently the slight biomagnification of heavy metals mercury (Hg) and lead (Pb) in fish tissues collected from the coastal marine areas of Tanzania. Therefore, a regular monitoring of heavy metal levels in fishes is necessary. Despite these observations, heavy metal levels in fish muscles in this study were generally below the permissible limits suggested by FAO [36] and posed no threat to public health yet.

Reference

- [1] Kravitz, M., Kroner, S., and Boethling, R. 2000. *Bioaccumulation Testing and Interpretation for the Purpose of Sediment Quality Assessment: Status and Needs*. Report no: EPA-823-R-00-001, US Environmental Protection Agency (EPA), Washington.
- [2] Mackay, D., and Fraser, A. 2000. "Bioaccumulation of Persistent Organic Chemicals: Mechanisms and Models." *Environmental Pollution* 110 (3): 375-91.
- [3] Connolly, J. P., and Pedersen, C. J. 1988. "A Thermodynamic-based Evaluation of Organic Chemical Accumulation in Aquatic Organisms." *Environmental Science & Technology* 22 (1): 99-103.
- [4] Rand, G. M., Wells, P. G., and McCarty, L. S. 1995. "Introduction to Aquatic Toxicology." In *Fundamentals of Aquatic Toxicology Effects, Environmental Fate, and Risk Assessment*. Taylor and Francis Publishers, North Palm Beach, Florida, USA, 3-67.
- [5] Gray, J. S. 2002. "Biomagnification in Marine Systems: The Perspective of an Ecologist." *Marine Pollution Bulletin* 45 (1): 46-52.
- [6] Gobas, F. A. 2000. "Bioconcentration and Biomagnification in the Aquatic Environment." In *Handbook of Property Estimation Methods for Chemicals, Environmental and Health Sciences*, edited by Boethling, R.S., and Mackay, D. Lewis, Boca Raton, 189-231.
- [7] Spacie, A., McCarty, L. S., and Rand, G. M. 1995. "Bioaccumulation and Bioavailability in Multiphase Systems." In *Fundamentals of Aquatic Toxicology* (2nd Edition), edited by Rand G. M. Washington, DC: Taylor and Francis, 493-521.
- [8] Barber, M. C., Suarez, L. A., and Lassiter, R. R. 1991. "Modelling Bioaccumulation of Organic Pollutants in Fish with an Application to PCBs in Lake Ontario Salmonids." *Canadian Journal of Fisheries & Aquatic Sciences* 48 (2): 318-37.
- [9] Wang, W. X. 2002. "Interactions of Trace Metals and Different Marine Food Chains" *Marine Ecology Progress Series* 243: 295-309.
- [10] Rounick, J. S., Winterbourn, M. J., and Lyon, G. L. 1982. "Differential Utilization of Allochthonous and Autochthonous Inputs by Aquatic Invertebrates in Some New Zealand Streams: A Stable Carbon Isotope Study." *Oikos*: 191-8.
- [11] Hamilton, S. K., Lewis, W. M., and Sippel, S. J. 1992. "Energy Sources for Aquatic Animals in the Orinoco River Floodplain: Evidence from Stable Isotopes." *Oecologia* 89 (3): 324-30.
- [12] Atwell, L., Hobson, K. A., and Welch, H. E. 1998. "Biomagnification and Bioaccumulation of Mercury in an Arctic Marine Food Web: Insights from Stable Nitrogen Isotope Analysis." *Canadian Journal of Fisheries & Aquatic Sciences* 55 (5): 1114-21.
- [13] Thompson, D. R., Furness, R. W., and Monteiro, L. R. 1998. "Seabirds as Biomonitorers of Mercury Inputs to Epipelagic and Mesopelagic Marine Food Chains." *Science of the Total Environment* 213 (1): 299-305.
- [14] Power, M., Klein, G. M., Guiguer, K., and Kwan, M. 2002. "Mercury Accumulation in the Fish Community of a Sub-Arctic Lake in Relation to Trophic Position and Carbon Sources." *Journal of Applied Ecology* 39 (5): 819-30.
- [15] Campbell, L. M., Norstrom, R. J., Hobson, K. A., Muir, D. C., Backus, S., and Fisk, A. T. 2005. "Mercury and Other Trace Elements in a Pelagic Arctic Marine Food Web (Northwater Polynya, Baffin Bay)." *Science of the Total Environment* 351: 247-63.
- [16] Luoma, S. N., and Rainbow, P. S. 2008. *Metal Contamination in Aquatic Environments: Science and Lateral Management*. Cambridge University Press.
- [17] Machiwa, J. F. 2000. "Heavy Metals and Organic Pollutants in Sediments of Dar es Salaam Harbour Prior to Dredging in 1999." *Tanzania Journal of Science* 26 (1): 29-46.
- [18] Kotze, P., Du Preez, H. H., and Van Vuren, J. H. J. 1999. "Bioaccumulation of Copper and Zinc in Oreochromis Mossambicus and Clarias Gariepinus, from the Olifants River, Mpumalanga, South Africa." *Water South Africa-Pretoria* 25: 99-110.
- [19] Ikemoto, T., Tu, N. P. C., Okuda, N., Iwata, A., Omori, K., Tanabe, S., et al. 2008. "Biomagnification of Trace Elements in the Aquatic Food Web in the Mekong Delta, South Vietnam Using Stable Carbon and Nitrogen Isotope Analysis." *Archives of Environmental Contamination & Toxicology* 54 (3): 504-15.
- [20] Revenga, J. E., Campbell, L. M., Arribere, M. A., and Guevara, S. R. 2012. "Arsenic, Cobalt and Chromium Food Web Biodilution in a Patagonia Mountain Lake." *Ecotoxicology & Environmental Safety* 81: 1-10.

- [21] France, R., and Steedman, R. 1996. "Energy Provenance for Juvenile Lake Trout in Small Canadian Shield Lakes as Shown by Stable Isotopes." *Transactions of the American Fisheries Society* 125 (4): 512-8.
- [22] Newell, R. I. E., Marshall, N., Sasekumar, A., and Chong, V. C. 1995. "Relative Importance of Benthic Microalgae, Phytoplankton, and Mangroves as Sources of Nutrition for Penaeid Prawns and Other Coastal Invertebrates from Malaysia." *Marine Biology* 123 (3): 595-606.
- [23] Bouillon, S., Koedam, N., Raman, A., and Dehairs, F. 2002. "Primary Producers Sustaining Macro-invertebrate Communities in Intertidal Mangrove Forests." *Oecologia* 130 (3): 441-8.
- [24] Grebmeier, J. M., McRoy, C. P., and Feder, H. M. 1988. "Pelagic-benthic Coupling on the Shelf of the Northern Bering and Chukchi Seas. I. Food Supply Source and Benthic Biomass." *Marine Ecology Progress Series* 48 (1): 57-67.
- [25] Achituv, Y., Brickner, I., and Erez, J. 1997. "Stable Carbon Isotope Ratios in Red Sea Barnacles (Cirripedia) as an Indicator of Their Food Source." *Marine Biology* 130 (2): 243-7.
- [26] Ponsard, S., and Arditi, R. 2000. "What Can Stable Isotopes ($\delta^{15}\text{N}$ and $\delta^{13}\text{C}$) Tell about the Food Web of Soil Macro-invertebrates?" *Ecology* 81 (3): 852-64.
- [27] Xu, J., Xie, P., Zhang, M., and Yang, H. 2005. "Variation in Stable Isotope Signatures of Seston and a Zooplanktivorous Fish in a Eutrophic Chinese Lake." *Hydrobiologia* 541 (1): 215-20.
- [28] Sheaves, M., and Molony, B. 2000. "Short-circuit in the Mangrove Food Chain." *Marine Ecology Progress Series* 199: 97-109.
- [29] Alfaro, A. C., Thomas, F., Sargent, L., and Duxbury, M. 2006. "Identification of Trophic Interactions within an Estuarine Food Web (Northern New Zealand) Using Fatty Acid Biomarkers and Stable Isotopes." *Estuarine, Coastal and Shelf Science* 70 (1): 271-86.
- [30] Cabana, G., and Rasmussen, J. B. 1994. "Modelling Food Chain Structure and Contaminant Bioaccumulation Using Stable Nitrogen Isotopes." *Nature* 372 (6503): 255-7.
- [31] Bowles, K. C., Apte, S. C., Maher, W. A., Kawei, M., and Smith, R. 2001. "Bioaccumulation and Biomagnification of Mercury in Lake Murray, Papua New Guinea." *Can. J. Fish. Aquat. Sci.* 58 (5): 888-97.
- [32] Mason, R. P., Laporte, J. M., and Andres, S. 2000. "Factors Controlling the Bioaccumulation of Mercury, Methylmercury, Arsenic, Selenium, and Cadmium by Freshwater Invertebrates and Fish." *Archives of Environmental Contamination & Toxicology* 38 (3): 283-97.
- [33] Richmond, M. D. 1997. *A Guide to the Seashores of Eastern Africa and the Western Indian Ocean Islands*. SIDA (Swedish International Development Cooperation Agency).
- [34] Bianchi, G. 1985. *FAO Species Identification Sheets for Fishery Purposes. Field Guide to the Commercial Marine and Brackish-water Species of Tanzania*. Pakistan: Fisheries and Aquaculture Department.
- [35] Newman, M. C., and Zhao, Y. 2005. "Measuring Metals and Metalloids in Water, Sediment and Biological Tissues." In *Techniques in Aquatic Toxicology* (2nd Edition), edited by Ostrander G. K. Boca Raton, FL: Lewis Publishers/CRC Press, 591-615.
- [36] Nauen, C. E. 1983. "Compilation of Legal Limits for Hazardous Substances in Fish and Fisheries Product." FAO, 102.
- [37] Cabana, G., and Rasmussen, J. B. 1996. "Comparison of Aquatic Food Chains Using Nitrogen Isotopes." *Proceedings of the National Academy of Science* 93 (20): 10844-7.
- [38] Lopez-Fernandez, H., and Winemiller, K. O. 2005. "Status of *Dionda Diaboli* and Report of Established Populations of Exotic Fish Species in Lower San Felipe Creek, Val Verde County, Texas." *The Southwestern Naturalist* 50 (2): 246-51.
- [39] Vander-Zanden, M. J., Cabana, G., and Rasmussen, J. B. 1997. "Comparing Trophic Position of Freshwater Fish Calculated Using Stable Nitrogen Isotope Ratios ($\delta^{15}\text{N}$) and Literature Dietary Data." *Canadian Journal of Fisheries & Aquatic Sciences* 54 (5): 1142-58.
- [40] Merritt, R. W., and Cummins, K. W., eds. 1996. *An Introduction to the Aquatic Insects of North America*. Kendall Hunt.
- [41] Jennings, S., Pinnegar, J. K., Polunin, N. V., and Warr, K. J. 2002. "Linking Size-based and Trophic Analyses of Benthic Community Structure." *Marine Ecology Progress Series* 226: 77-85.
- [42] Vander-Zanden, M. J., and Rasmussen, J. B. 2001. "Variation in $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ Trophic Fractionation: Implications for Aquatic Food Web Studies." *Limnology & Oceanography* 46 (8): 2061-6.
- [43] Thomson, D., Maher, W., and Foster, S. 2007. "Arsenic and Selected Elements in Inter-tidal and Estuarine Marine Algae, South-east Coast, NSW, Australia." *Applied Organometallic Chemistry* 21 (6): 396-411.
- [44] Rahman, M. A., Hasegawa, H., and Lim, R. P. 2012. "Bioaccumulation, Biotransformation and Trophic Transfer of Arsenic in the Aquatic Food Chain." *Environmental Research* 116: 118-35.
- [45] Barwick, M., and Maher, W. 2003. "Biotransference and Biomagnification of Selenium Copper, Cadmium, Zinc, Arsenic and Lead in a Temperate Seagrass Ecosystem from Lake Macquarie Estuary, NSW, Australia." *Marine Environmental Research* 56 (4): 471-502.

- [46] Hardersen, S., and Wratten, S. D. 1998. "The Effects of Carbaryl Exposure of the Penultimate Larval Instars of *Xathocnemis Zealandica* on Emergence and Fluctuating Asymmetry." *Ecotoxicology* 7 (5): 297-304.
- [47] Fernandes, T. F., Christofi, N., and Stone, V. 2007. "The Environmental Implications of Nanomaterials." In *Nanotoxicology: Characterization, Dosing and Health Effects*, edited by Monteiro-Riviere, N., and Tran, C. L. CRC Press, Boca Raton, FL, USA, 405-20.
- [48] Woolson, E. A. 1975. "Bioaccumulation of Arsenicals." In *Arsenical Pesticides*, edited by Woolson, E. A. Washington, D.C.: American Chemical Society, ACS Symposium.
- [49] Wagemann, R., Snow, N. B., Rosenberg, D. M., and Lutz, A. 1978. "Arsenic in Sediments, Water and Aquatic Biota from Lakes in the Vicinity of Yellowknife, Northwest Territories, Canada." *Archives of Environmental Contamination & Toxicology* 7 (1): 169-91.
- [50] Chen, C. Y., and Folt, C. L. 2000. "Bioaccumulation and Diminution of Arsenic and Lead in a Freshwater Food Web." *Environmental Science & Technology* 34 (18): 3878-84.
- [51] Karez, C. S., Magalhaes, V. F., Pfeiffer, W. C., and Amado Filho, G. M. 1994. "Trace Metal Accumulation by Algae in Sepetiba Bay, Brazil." *Environmental Pollution* 83 (3): 351-6.
- [52] Lee, B. G., Griscom, S. B., Lee, J. S., Choi, H. J., Koh, C. H., Luoma, S. N., et al. 2000. "Influences of Dietary Uptake and Reactive Sulfides on Metal Bioavailability from Aquatic Sediments." *Science* 287 (5451): 282-4.
- [53] Romeo, M., Siau, Y., Sidoumou, Z., and Gnassia-Barelli, M. 1999. "Heavy Metal Distribution in Different Fish Species from the Mauritania Coast." *Science of the Total Environment* 232 (3): 169-75.
- [54] Kidd, K. A., Schindler, D. W., Hesslein, R. H., and Muir, D. C. G. 1995. "Correlation between Stable Nitrogen Isotope Ratios and Concentrations of Organochlorines in Biota from a Freshwater Food Web." *Science of the Total Environment* 160: 381-90.
- [55] Al-Reasi, H. A., Ababneh, F. A., and Lean, D. R. 2007. "Evaluating Mercury Biomagnification in Fish from a Tropical Marine Environment Using Stable Isotopes ($\delta^{13}\text{C}$ and $\delta^{15}\text{N}$)." *Environmental Toxicology & Chemistry* 26 (8): 1572-81.
- [56] Luoma, S. N. 1989. "Can We Determine the Biological Availability of Sediment-bound Trace Elements?" *Hydrobiologia* 176 (1): 379-96.
- [57] Mamboya, F. A. 2007. "Heavy Metal Contamination and Toxicity: Studies of Macroalgae from the Tanzania Coast." Doctoral dissertation, Botaniska institutionen.
- [58] Elmaci, A., Teksoy, A., Topac, F. O., and Özenin, N. 2007. "Assessment of Heavy Metals in Lake Uluabat, Turkey." *African Journal of Biotechnology* 6 (19): 2236-44.
- [59] Eisler, R. 1981. *Trace Metal Concentrations in Marine Organisms*. New York: Pergamon Press.
- [60] Rainbow, P. S. 1993. "The Significance of Trace Metal Concentrations in Marine Invertebrates." In *Ecotoxicology of Metals in Invertebrates*, edited by Dallinger, R., and Rainbow, P. S. Chelsea, USA: Lewis Publishers, 3-23.
- [61] Romeo, M., Siau, Y., Sidoumou, Z., and Gnassia-Barelli, M. 1999. "Heavy Metal Distribution in Different Fish Species from the Mauritania Coast." *Science of the Total Environment* 232 (3): 169-75.
- [62] Canli, M., and Atli, G. 2003. "The Relationships between Heavy Metal (Cd, Cr, Cu, Fe, Pb, Zn) Levels and the Size of Six Mediterranean Fish Species." *Environmental Pollution* 121 (1): 129-36.
- [63] Friberg, L., Norberg, G. F., and Vouk, V. B. 1979. *Handbook of Metal Toxicology*. Amsterdam: Elsevier/North Holland Biomedical Press.