Screening for Sodium-Salt Tolerance at the Seedling Stage of Rice (*Oryza sativa* L.) Genotypes from Sudan and South Sudan

Seif Gasim¹, Ishraka Abuanja¹, Mohamedsalih Dafalla² and Abdelwahab Abdalla¹

1. Department of Agronomy, Faculty of Agriculture, University of Khartoum, Shambat 13314, Sudan

2. Department of Soil Science, Faculty of Agriculture, University of Khartoum, Shambat 13314, Sudan

Abstract: Morpho-physiological and genetic studies of salinity are important in understanding the mechanism of plant adaptation to stressful environment. Eighteen rice genotypes collected from Sudan and South Sudan, which were never tested for salt tolerance, as well as two genotypes (*FL478* as tolerant check and *IR29* as sensitive check) from the International Rice Research Institute (IRRI), were subjected to salinity stress at seedling stage. Test was carried out in hydroponic system applying electrical conductivity (EC) 12 dS/m NaCl using randomized complete block design with three replicates. Most of the genotypes showed sensitivity to salt stress; one genotype *PIPANFARY RED2* was moderately sensitive and three genotypes *MASURY1*, *MASURY2* and *FL478* were tolerant. Salinity significantly reduced leaf dry weight, shoot dry weight, root dry weight and biomass production (biomass/plant) by 31%, 42%, 60% and 47%, respectively. Tolerant genotypes accumulated low amount (2.52 g/100 g dry weight (dwt)) of Na⁺ in the root, whereas highly sensitive genotypes accumulated high amount (3.87 g/100 g dwt) of Na⁺. Tolerant genotypes showed less reduction in K⁺ concentration than the sensitive genotypes. Therefore, they maintained lowest Na⁺/K⁺ ratio in the shoot (1.47%) than in the root (3.69%) compared to the intolerant genotypes (7.49% and 8.49%). The genotypes that showed tolerance to salinity stress can be used as a source of resistance/tolerance in a breeding program for rice improvement in uplands areas in semi-arid condition.

Key words: Rice, salt tolerance, seedling stage, plant tissue.

1. Introduction

Salinity is a major constraint restricting rice productivity across many rice production areas [1]. It is a worsening problem particularly in inland areas of arid and semi-arid due to the buildup of salinity as a consequence of excessive use of irrigation water with improper drainage coupled with the use of poor quality irrigation.

Rice susceptibility to salinity changes with the stage of development; germination, active tillering and maturation stages are less susceptible to salinity than to the seedling and early reproductive stages [1, 2]. Recently, considerable progress was made in understanding the physiological bases of salt tolerance

in rice and salt tolerant genotypes were developed [3, 4]. However, due to their local adaptation, landraces cultivars are being used as potential genetic resources of stress tolerance traits [5].

A large contributor to salinity stress is the build-up of the sodium ion (Na⁺) in the cytoplasm of leaf cells. High cytoplasmic Na⁺ interferes with processes that require the binding of potassiumion (K⁺), protein synthesis and the activation of key metabolic enzymes [6]. One key mechanism of salinity tolerance is the ability of a plant to control Na⁺ transport at both the tissue and cell levels, either by secreting Na⁺ into tissues, cells and organelles where it can do little damage, or by minimizing the amount of Na⁺ entering the plant through its roots [6, 7]. In Sudan, most of the rice cultivated areas are of flood and salinity. However, no information is available on the

Corresponding author: Seif Gasim, associate professor, research field: plant breeding.

mechanism of salinity tolerance of the locally grown cultivars. Therefore, the objective of the present study was to evaluate salinity tolerance of some Sudanese genotypes (never evaluated for their salt tolerance) during the early seedling stage and to identify the mechanism of tolerance with reference to Na^+ and K^+ uptake.

2. Materials and Methods

2.1 Planting Material, Experimental Design and Evaluation of Salinity

Eighteen rice genotypes were studied: seven of them were collected from Aweil, South Sudan (*NBGS1*, *NBGS2*, *NBGS3*, *NBGA*, *BG400-1*, *BG90-2*, *BANBAN*) and eleven from Kosti, Sudan (*MASURY1*, *PIPANFARY RED1*, *JAING ARRI RED*, *SOMMBOY*, *COMARWA*, *PAINJLA*, *PIPANFARY*, *TAGMIZEDO*, *PIPANFARY RED2*, *BACTING ARRI*, *MASURY2*). In addition, two genotypes were provided by the International Rice Research Institute (IRRI) namely *IR29* and *FL478* used for salinity check; *IR29* as sensitive check and *FL478* as tolerant check [8, 9]. The seeds were first incubated for 3 d in an oven at 50 °C to break seed dormancy, soaked in distilled water and then incubated and pre-germinated in the dark at 32 °C until the seeds sprouted (in about 48 h). Five germinating seeds from each genotype were transferred to styrofoam floats in a plastic container of 8 L capacity filled with deionized water and the entire setup was transferred to the green house of the IRRI. A randomized complete block design was used to carry out the experiment; entries were replicated three times. Two sets of treatments were used: one normal and the other saline. Three days later, the deionized water was replaced by Yoshida nutrient solution [10] for 4 d. Electrical conductivity (EC) of the solution was adjusted by the addition of NaCl into 6 dS/m after 3 d and 12 dS/m after 6 d. The pH was checked and adjusted every day within the range of 5.0-5.4 and nutrient solution was changed every week to maintain nutrients in the solution. IRRI's Standard Evaluation

System (SES) for rice [11] was used for evaluation of salinity tolerance in score (3 = tolerant; 5 = moderate tolerant; 7 = susceptible; 9 = sensitive). Salinity effect started to appear in some genotypes from the second scoring then most of them reached high scoring of 7-9 after 14 d from subjecting the plants to salinity.

2.2 Na^+ and K^+ Determination

The plant samples (shoots and roots, separately) were transferred into labeled coin envelopes and put in an oven for drying at 70 °C to a constant weight. Then their dry weights were determined and homogenized using scissor and beaker and sub-samples of 20.5 mg each were introduced into labeled 50 mL Falcon tubes. Then 20 mL 0.1 M of acetic acid was added to each tube using an automatic dispensing burette. Extreme care was taken to thoroughly wet the sample with the acid, and the samples were digested in an oven at 60 °C for overnight per a couple of 5 d, then the bottles were shaken gently and filtered through a Whatman filter papers No.1 into new set of labeled 50 mL Falcon tubes, and finally 1 mL of the filtered sample was transferred to 14 mL Falcon tubes using pipette, and diluted to 10 mL using Nano pure water for Na^+ and K^+ determination on an atomic absorption spectrometer (AAS) (Perkin Elmer A Analyst 400). Then Na^+ and K^+ concentrations were determined in percentage as follows:

$$\% \text{Na}^+ \text{ or } \text{K}^+ = \frac{\text{ppm volume} \left(\frac{20}{1000} \right) \times 100}{\text{Sample dry weight}} \quad (1)$$

where, ppm volume is the amount of Na^+ or K^+ from the AAS, 20/1000 is the amount of solvent used for extraction in liter and 10 is the diluted volume. Moreover, leaf dry weight, shoot dry weight, root dry weight and biomass/plant were recorded under saline and control conditions.

2.3 Screening for Na^+ Tolerance in Plant Tissue and Leaf to Leaf Compartmentation

Based on the results of the effect of the salinity in the studied genotypes, two tolerant genotypes

(*MASURY1* and *MASURY2*) and two sensitive genotypes (*BG400-1* and *SOMMBOY*) were selected, screened and compared with *FL478* (tolerant check) and *IR29* (sensitive check) for shoot sodium, tissue Na^+ tolerance and leaf to leaf compartmentation. The experiment was laid out using different salinity treatments (EC0, EC6, EC12 and EC18 or 0, 60, 90 and 120 mM) with three replications in a randomized complete block design. Seed treatments, incubation, germination and salinity treatments were carried out following the procedure mentioned above (the trays designed for different salinity treatments were salinized after 14 d). Depending on the toxicity symptoms, the individual leaves (the youngest fully expanded leaf) of the genotype encompassing the range of observable symptoms in the highest salinity treatment (EC18) which reached an average SES of 9 were harvested from all salinity treatments and the control according to a wide range of damage that appeared. Then the samples were oven dried, weighed and Na^+ concentrations were determined as above. For leaf to leaf compartmentation screening, the seedlings were left to grow under Yoshida nutrients solution for 21 d then salinized with NaCl (6 g NaCl/L; 12 dS/m). In the following day, the leaf three (L3) and leaf four (L4) from two plants of each genotype were harvested (4 d after salinization). The samples were oven dried and weighed and subsamples were taken for Na^+ extraction, and Na^+ concentration was determined on an AAS as mentioned before.

2.4 Statistical Analysis

Analysis of variance was carried out on the collected data using software Statistical Tool for Agricultural Research [12]. Least significant difference ($\text{LSD}_{0.05}$) was used for mean separation.

3. Results

Statistical analysis (data not shown) revealed that the effect of genotype and salt concentration level (salinity) were highly significant ($p \leq 0.01$) for all of

the phenotypic parameters under study (leaf dry weight, shoot dry weight, root dry weight and biomass/plant) and physiological traits (shoot Na^+ , shoot K^+ , shoot Na^+/K^+ ratio, root Na^+ , root K^+ and root Na^+/K^+ ratio). The effects of genotype, salinity as well as the effect of genotype X salt concentration interaction were significant for all traits, except for root Na^+ , which revealed non-significant differences. When salt stress of 12 dS/m was applied, all of Aweil genotypes (South Sudan) were susceptible to salinity (SES values of 7.0-8.5). For Kosti genotypes (Sudan), two genotypes (*MASURY1* and *MASURY2*) showed tolerance to salinity (SES values of 3.0-4.5), one genotype (*PIPANFARY RED2*) was moderately tolerant (SES value of 5-6.7) and one genotype (*SOMMBOY*) was highly sensitive to salinity stress (SES value of 9); the rest of the genotypes were susceptible (SES of 7.0-8.5) as shown in Table 1. Salinity caused great reduction in phenotypic parameters (leaf dry weight, shoot dry weight, root dry weight and biomass production) in all genotypes, except for *FL478* (tolerant check), which showed an increase in shoot dry weight by 12% and in biomass/plant by 8% under saline condition (Table 1). On average, salinity caused a considerable reduction in leaf dry weight, shoot dry weight, root dry weight and biomass production by 31%, 42%, 60% and 47%, respectively, after 14 d from salinization with high variation among genotypes. The results also showed that roots dry weight was the most severely affected trait under saline condition.

Table 2 shows the variation in the percentage of Na^+ and K^+ in shoots and roots of the genotypes measured after 14 d from salinization. Na^+ concentration of different plant tissues of all genotypes was increased under saline condition than under control condition. The percentage of Na^+ concentration in the shoot ranged from 2.354% (*MASURY2*) to 4.874% (*BANBAN*) and ranged in the root from 1.899% (*NBGS3*) to 3.285% (*TAGMIZEDO*). However, tolerant genotypes accumulated less amount

**Screening for Sodium-Salt Tolerance at the Seedling Stage of Rice (*Oryza sativa* L.)
Genotypes from Sudan and South Sudan**

Table 1 Variation in 20 rice genotypes in Standard Evaluation System (SES) score, leaf dry weight, shoot dry weight, root dry weight and biomass/plant after 14 d from salinization under saline and control conditions.

Genotype	Saline	Control	Saline	Control	Saline	Control	Saline	Control	Saline	Control
	SES score	SES score	Leaf dry weight (mg)	Leaf dry weight (mg)	Shoot dry weight (mg)	Shoot dry weight (mg)	Root dry weight (mg)	Root dry weight (mg)	Biomass /plant (mg)	Biomass /plant (mg)
<i>NBGS1</i>	7.7	1.0	17.1	26.9	278.2	480.5	74.2	154.4	176.2	317.5
<i>NBGS2</i>	7.7	1.0	21.3	26.7	312.2	508.8	78.9	154.7	195.6	331.8
<i>NBGS3</i>	8.0	1.0	18.9	40.9	221.1	607.8	60.3	217.5	140.7	412.7
<i>NBGA</i>	7.0	1.0	22.0	37.7	300.9	560.0	78.3	229.8	189.6	394.9
<i>BG400-1</i>	8.3	1.0	10.5	15.8	200.6	401.1	55.8	147.6	128.2	274.4
<i>BG90-2</i>	7.0	1.0	22.0	31.0	336.6	738.3	75.7	242.5	206.1	490.4
<i>BANBAN</i>	8.0	1.0	32.4	47.9	243.1	690.3	61.9	263.5	152.5	476.9
<i>MASURY1</i>	4.3	1.0	33.7	53.5	653.4	748.3	175.4	261.0	414.4	504.6
<i>PIPANFARY RED1</i>	8.3	1.0	25.6	29.0	335.4	746.7	101.7	348.9	218.6	547.8
<i>JAING ARRI RED</i>	7.7	1.0	25.3	30.9	382.8	592.5	115.6	301.9	249.2	447.2
<i>SOMMBOY</i>	9.0	1.0	33.1	57.1	368.2	862.9	96.7	480.8	232.5	671.8
<i>COMARWA</i>	7.7	1.0	18.8	27.3	439.6	739.2	152.9	356.2	296.3	547.7
<i>PAINJLA</i>	8.3	1.0	33.7	30.0	356.8	683.2	107.1	355.9	232.0	519.5
<i>PIPANFARY</i>	8.3	1.0	17.9	34.0	296.4	465.3	93.2	267.1	194.8	366.2
<i>TAGMIZEDO</i>	7.0	1.0	28.6	36.4	542.5	907.1	175.5	427.3	359.0	667.2
<i>PIPANFARY RED2</i>	5.3	1.0	28.5	37.9	476.8	629.6	158.4	269.8	317.6	449.7
<i>BACTING ARRI</i>	7.0	1.0	20.9	30.5	397.8	668.3	108.1	371.4	253.0	519.9
<i>MASURY2</i>	3.7	1.0	26.1	41.3	511.0	655.0	159.3	249.8	335.1	452.4
<i>FL478</i> (tolerant check)	3.0	1.0	20.3	28.2	634.6	564.7	170.9	178.3	402.8	371.5
<i>IR29</i> (sensitive check)	9.0	1.0	13.4	16.6	204.9	428.0	66.8	152.5	135.8	290.2
Mean	7.1	1.0	23.5	33.9	367.5	633.9	108.34	271.5	241.5	452.7
LSD _{0.05}	1.3		4.1		79.2		44.9		58.7	

of Na⁺ in the shoot with a mean value of 2.52 g/100 g dry weight (dwt), whereas susceptible genotypes accumulated higher amounts of Na⁺ with a mean value of 3.87 g/100 g dwt. On the other hand, the tolerant genotypes accumulated highest amount of Na⁺ in the root (2.51 g/100 g dwt) than the highly sensitive genotypes (2.44 g/100 g dwt) (Table 2).

Nevertheless, sensitive genotypes absorbed more Na⁺ and transferred high amount to the shoot, whereas the tolerant genotypes absorbed less Na⁺ and retained most of it in their roots, therefore maintained higher shoot growth rate (Table 2). Due to the antagonistic properties of Na⁺ and K⁺ with increase in salt concentration, K⁺ rate was decreased in all tissues, with variations among genotypes, except for the tolerant genotype *MASURY1*, which showed an increase in K⁺ concentration in the shoot by 3% (Table 2). The K⁺ concentration in the shoot ranged

from 0.545% (*IR29* (sensitive check)) to 2.005% (*MASURY1*) and in the root it ranged from 0.181% (*PIPANFARY RED1*) to 0.806% (*MASURY2*).

The results in Table 3 show that salinity caused an increase in the ratio of Na⁺/K⁺ in both shoot and root. The highest ratio of Na⁺/K⁺ in the shoot was related to highly sensitive genotype (*SOMMBOY*, 7.499%), whereas the lowest ratio was related to tolerant genotype (*MASURY1*, 1.306%). On the other hand, the highest ratio of Na⁺/K⁺ in root was related to highly sensitive genotype (*JAING ARRI RED*, 13.951%), while the lowest ratio was related to moderately tolerant genotype (*MASURY2*, 3.211%). The shoot and the root Na⁺/K⁺ ratios increased under saline condition by 98% and 92% under control condition. Under saline condition, the tolerant genotypes exhibited lower amounts of Na⁺/K⁺ ratio in the shoot (1.468%) than in the root (3.690%). But the highly

Table 2 Variation in SES score and Na⁺ and K⁺ concentrations in shoots and roots of 20 rice genotypes measured after 14 d from salinization under saline and control conditions.

Genotype/source	Shoot		Root		Shoot		Root	
	Saline	Control	Saline	Control	Saline	Control	Saline	Control
	Na ⁺ content	Na ⁺ content	Na ⁺ content	Na ⁺ content	K ⁺ content	K ⁺ content	K ⁺ content	K ⁺ content
<i>NBGS1</i>	3.370	0.036	2.587	0.515	1.041	2.572	0.454	0.908
<i>NBGS2</i>	3.140	0.048	2.756	0.539	1.293	2.646	0.694	1.122
<i>NBGS3</i>	3.320	0.029	2.899	0.599	0.937	1.880	0.361	1.003
<i>NBGA</i>	3.499	0.053	2.762	0.681	1.264	2.022	0.540	0.968
<i>BG400-1</i>	3.125	0.087	2.441	0.356	0.862	1.555	0.389	0.855
<i>BG90-2</i>	3.982	0.058	2.420	0.645	0.863	2.663	0.385	0.988
<i>BANBAN</i>	4.874	0.103	2.805	0.602	0.921	1.963	0.342	1.010
<i>MASURY1</i>	2.609	0.043	2.445	0.457	2.005	1.955	0.556	0.828
<i>PIPANFARY RED1</i>	4.315	0.141	2.268	0.536	0.771	2.112	0.181	0.857
<i>JAING ARRI RED</i>	3.936	0.156	2.572	0.526	0.686	1.917	0.185	0.840
<i>SOMMBOY</i>	3.761	0.155	2.820	0.622	0.571	2.045	0.229	0.918
<i>COMARWA</i>	3.753	0.233	2.921	0.566	0.815	1.885	0.235	0.932
<i>PAINJLA</i>	3.563	0.141	2.822	0.424	0.670	1.817	0.316	1.085
<i>PIPANFARY</i>	4.091	0.262	2.752	0.687	0.607	1.982	0.299	0.942
<i>TAGMIZEDO</i>	3.769	0.137	2.285	0.603	0.997	1.905	0.366	0.936
<i>PIPANFARY RED2</i>	2.429	0.067	2.715	0.555	1.418	1.895	0.677	0.999
<i>BACTING ARRI</i>	4.010	0.102	2.021	0.588	0.859	1.881	0.344	0.903
<i>MASURY2</i>	2.354	0.104	2.508	0.470	1.453	1.919	0.806	1.144
<i>FL478</i> (tolerant check)	2.607	0.094	2.583	0.604	1.802	2.129	0.757	0.890
<i>IR29</i> (sensitive check)	3.544	0.082	2.344	0.537	0.545	1.869	0.303	1.162
Mean	3.50	0.107	2.59	0.557	1.019	2.031	0.421	0.965
LSD _{0.05}	0.70		0.53		0.25		0.12	

sensitive genotypes showed highest amount of Na⁺/K⁺ ratio in the shoot (7.049) and root (10.133%). Under saline condition, tolerant genotypes maintained lowest Na⁺ concentrations, highest K⁺ concentrations and lowest Na⁺/K⁺ ratios in the shoot than in the root; whereas sensitive genotypes attained highest Na⁺ concentrations, lowest K⁺ concentrations, highest Na⁺/K⁺ ratios in the shoot than in the root (Tables 2 and 3).

Screening for Na⁺ in the selected accessions at different EC (0, 6, 12 and 18 dS/m) indicated that the effect of genotype, salinity levels and genotype × salinity interaction were highly significant ($p \leq 0.01$) in all plant leaves and in plant leaf tissue tolerance (data not shown). As a result, the mean values of genotypes and salinity levels were different from each other. In terms of tissue tolerance, tissue Na⁺ concentration increased with increasing the amount of

NaCl concentration; therefore, the highest level of Na⁺ concentration was reached with the level of EC18 (Table 4). The highest level of Na⁺ concentration was shown by *MASURY2* (4.178 g/100 g dwt), and the lowest level was obtained for the genotype *MASURY1* (2.183 g/100 g dwt). Compared to the tolerant check (*FL478*), *MASURY2* exhibited the highest Na⁺ concentration (3.446 g/100 g dwt), so the concentration of Na⁺ in the tissue of *MASURY2* was higher than that in the tolerant check. Moreover, the results showed that sensitive genotypes to salinity accumulated high Na⁺ concentration but scored low SES, whereas tolerant genotypes accumulated low Na⁺ and scored high SES (Fig. 1).

For leaf to leaf compartmentation, leaf three and leaf four, the amount of Na⁺ content increased from day to day and reached the highest level in day four (four days after salinization (DAS)). The highest amount

Table 3 Variation in Na⁺/K⁺ ratio in shoots and roots of 20 rice genotypes measured after 14 d from salinization.

Genotype	Shoot		Root	
	Saline	Control	Saline	Control
	Na ⁺ /K ⁺ ratio	Na ⁺ /K ⁺ ratio	Na ⁺ /K ⁺ ratio	Na ⁺ /K ⁺ ratio
<i>NBGS1</i>	3.547	0.015	6.207	0.560
<i>NBGS2</i>	2.722	0.020	4.601	0.486
<i>NBGS3</i>	4.278	0.015	7.593	0.596
<i>NBGA</i>	2.930	0.025	5.611	0.698
<i>BG400-1</i>	4.926	0.060	9.035	0.416
<i>BG90-2</i>	4.692	0.023	6.798	0.657
<i>BANBAN</i>	5.523	0.057	8.236	0.608
<i>MASURY1</i>	1.306	0.023	4.398	0.554
<i>PIPANFARY RED1</i>	5.797	0.068	12.298	0.627
<i>JAING ARRI RED</i>	6.489	0.087	13.951	0.627
<i>SOMMBOY</i>	7.499	0.077	12.368	0.683
<i>COMARWA</i>	6.298	0.124	7.823	0.608
<i>PAINJLA</i>	5.478	0.077	9.405	0.427
<i>PIPANFARY</i>	6.790	0.137	9.496	0.731
<i>TAGMIZEDO</i>	3.835	0.072	9.365	0.660
<i>PIPANFARY RED2</i>	1.981	0.035	4.213	0.559
<i>BACTING ARRI</i>	4.899	0.055	9.788	0.662
<i>MASURY2</i>	1.652	0.053	3.211	0.414
<i>FL478</i> (tolerant check)	1.447	0.043	3.461	0.676
<i>IR29</i> (sensitive check)	6.600	0.045	7.898	0.463
Mean	4.434	0.056	7.788	0.586
LSD _{0.05}	1.03		1.68	

Table 4 Tissue sodium concentrations of six rice genotypes subjected to four different levels of salinity in hydroponic solution.

Accessions	Tissue sodium concentrations			
	EC0	EC6	EC12	EC18
<i>MASURY1</i>	0.016	0.100	0.125	2.183
<i>MASURY2</i>	0.007	0.056	0.929	4.178
<i>BG400-1</i>	0.007	0.039	0.818	2.954
<i>SOMMBOY</i>	0.028	0.330	1.615	3.714
<i>FL478</i> (tolerant check)	0.026	0.062	0.558	3.970
<i>IR29</i> (sensitive check)	0.028	0.144	1.199	3.446
Mean	0.112	0.180	0.870	3.41
LSD _{0.05}	0.003	0.012	0.002	0.001

(625.1 mg/100 g dwt) of Na⁺ content was scored by the genotype *SOMMBOY* and the lowest amount (30.7 mg/100 g dwt) was obtained for *BG400-1* (Table 5). On the other hand, sodium concentration in leaf three was higher than in leaf four for all rice accessions. The result also showed a negative correlation between leaf to leaf Na⁺ concentration and SES score; thus, the

genotypes with the highest leaf to leaf Na⁺ concentration gave the lowest SES score (Fig. 2). Similarly, as shown in Fig. 3, the genotypes with the lowest SES score under salinity condition exhibited the highest tissue tolerance score, higher leaf to leaf compartmentation score and lower shoot sodium concentration, and the reverse is true for genotypes with lowest SES score.

Fig. 1 Correlation between tissue tolerance score and shoot Na⁺ concentration in six rice genotypes.

Table 5 Variation in sodium concentration in leaf three (L3) and leaf four (L4) of six rice accessions measured in four successive days after salinization (DAS).

Accessions	Sodium concentration in L3				Sodium concentration in L4			
	1 DAS	2 DAS	3 DAS	4 DAS	1 DAS	2 DAS	3 DAS	4 DAS
MASURY1	30.7	84.0	183.5	318.2	1.0	1.0	9.3	86.9
MASURY2	12.0	200.6	245.7	367.9	1.3	1.4	6.0	164.1
BG400-1	60.1	65.5	113.2	278.1	1.0	13.8	22.6	30.7
SOMMBOY	89.8	459.4	610.1	993.8	5.0	455.1	485.1	625.1
FL478 (tolerant check)	75.7	115.7	139.9	251.6	1.0	1.0	4.0	40.2
IR29 (sensitive check)	38.1	229.7	294.3	350.9	41.6	99.2	147.4	193.4
Mean	51.07	192.48	264.45	426.75	8.48	110.37	112.40	190.07
LSD _{0.05}	10.83	30.2	33.73	46.83	3.09	33.58	34.53	21.36

Fig. 2 Correlation between leaf to leaf Na⁺ concentration and leaf to leaf compartmentation in six rice genotypes.

**Screening for Sodium-Salt Tolerance at the Seedling Stage of Rice (*Oryza sativa* L.)
Genotypes from Sudan and South Sudan**

Fig. 3 SES score, tissue tolerance score, shoot Na⁺ concentration and leaf to leaf compartmentation score of six rice genotypes.

4. Discussion

Salt tolerance in rice is complex, and involves several physiological and adaptive mechanisms [13, 14]. The physiological bases of salt tolerance during early seedling stage are fairly well understood; key traits include high seedling vigor with a consequent dilution of salt concentration in sensitive tissues, selective ion uptake at the root level, compartmentation of harmful ions in structural and old tissues in old leaves and roots, and high tissue tolerance through compartmentation of salts in the apoplast.

Genetic variation in salt tolerance has been frequently documented in rice as well as in other crop species [9]. However, sodium is the major cation that accumulates in plant tissues and organs when salinity increases. With significant effect of salt concentration and genotype interaction, different genotypes show different responses to salt concentration [15]. The non-significant effect of Na⁺ concentration on root dry weight of the genotypes indicated that the difference between genotypes was not attributed to sodium

accumulation in the roots, because most of the salt concentration was accumulated in the shoots rather than in the roots. When the rice genotypes were exposed to salt stress of 12 dS/m, 15 genotypes from Sudan were sensitive, one (*PIPANFARY RED2*) was moderate and two (*MASURY1* and *MASURY2*) were tolerant to salinity, suggesting that evaluation at such level of salt concentration could be sufficient to detect accessions with reasonably high levels of tolerance of salt stress [8]. Also, this is an indication that there may be a gene governing the tolerance of these genotypes to salt stress, which is absent in the rest of the genotypes. This gene would be useful in environments with mild or moderate salt stress early in the season.

In the present study, tolerant genotypes showed less reduction percentage in phenotypic parameters, and intolerant genotypes exhibited high reduction percentage. The severe reduction in phenotypic parameters as a result of salinity stress indicates seedling stage is more sensitive to salinity stress [16, 17]. The increase in shoot dry weight and

biomass/plant shown by *FL478* was mostly due to their tolerance to salinity as reported by Moradi *et al.* [8]. Moreover, the severe effect of salinity on root dry weight could be attributed to the accumulation of most of the salts in the roots than in the shoots [18, 19]. The significant variation among the genotypes due to salt stress indicates that increase in Na^+ concentration reduces K^+ concentration either in the shoots or in the roots. Under condition of high salt concentration, Na^+ concentration was higher in the shoot than in the root, which led to reduction of K^+ concentration in the shoot and increase of Na^+/K^+ ratio and this is one of the tolerance mechanisms in grasses, which limits the transport of Na^+ into photosynthetic cells and the growing of meristem active tissue [5]. Therefore, tolerant genotypes accumulated less amount of Na^+ in the shoot, high amount of K^+ and less ratio of Na^+/K^+ , hence retained most of Na^+ content to the root, thus they maintained higher shoot growth rate. These findings agree with those of Peng and Ismail [14]. Moreover, the high concentration of Na^+ in the plant treated with NaCl suggests that the culture of the NaCl solution imposed ionic stress in the plant, in addition to the osmotic stress. This is further evidenced by the antagonistic effect of higher levels of Na^+ on K^+ uptake, which has been shown to occur in previous studies with high concentrations of Na^+ in nutrient or soil solution [20].

The selected genotypes showed uniform green color under control condition. Tolerant (*MASURY1* and *MASURY2*) and moderately tolerant (*BG400-1*) genotypes to salinity stress under salinized condition indicating their ability to tolerate Na^+ concentration as they showed less growth reduction in the studied parameters than intolerant (*SOMMBOY*) genotype, which agrees with the report of Castillo *et al.* [21]. When genotypes were exposed to different EC (0, 6, 12 and 18 dS/m), the highest Na^+ concentration was reached at the level of EC18. The highest level of Na^+ concentration exhibited by the *MASURY2* (tolerant genotype) indicates that this genotype has a

mechanism of compartmentation of salts in the apoplast to avoid buildup of toxic concentrations in the cytoplasm. This tolerance could probably be achieved through a set of active processes involving a gene family of ion transporters such as Na^+/H^+ antiporters that sequester salt in vacuoles [22] or move it out of the cell cytoplasm and recirculate it back to roots [23]. The highly significant variations between the genotypes for salt accumulation in leaf three (old leaf) than in leaf four (new leaf) suggesting the accumulation of salts in older leaves than in the newly growing leaves. The highest and lowest levels of Na^+ concentrations in leaf three and leaf four achieved by the sensitive genotype (*SOMMBOY*) and the moderately tolerant genotype (*BG400-1*) in day four indicate that this moderately tolerant genotype has a mechanism of tissue tolerance by accumulation of Na^+ in the vacuoles of the cell of the old leaves [6]. However, the lower SES score for the tolerant genotypes and the higher tissue tolerance score, higher leaf to leaf compartmentation score and lower shoot sodium concentration than for the intolerant genotypes indicate that genotypes with higher tissue tolerance maintain their cellular ultrastructure and have higher net photosynthesis at higher amounts of tissue salt concentration [3].

5. Conclusions

It could be concluded that salt stress severely affect the physiological traits, causing highly significant variability among the genotypes and the interaction between them. Tolerant genotypes accumulated lower amounts of sodium in the shoot and less reduction in K^+ concentration than the sensitive genotypes. Therefore, genotypes that showed tolerance to sodium salt stress could be used in breeding programs for tolerance of rice plants to salt stress in Sudan. Much work is needed to study the effect of salinity and sodicity on electrolyte concentration on growth parameters in rice plants in Sudan. In addition, understanding the molecular genetics and physiology

of the traits affected by salt stress will form the basis for further improvements in salt tolerance of crop species.

Acknowledgments

Thanks are due to the ministry of Higher Education and Scientific Research for providing the fund to carry out the research work and to IRRI technicians for the technical assistance. Thanks are also extended to the Agricultural Research Corporation, Kosti, Sudan and Ministry of Agriculture, Bar Elgabal state, South Sudan for their help in providing the germplasm.

References

- [1] Ali, M. N., Yeasmin, L., Gantait, S., Goswami, R., and Chakrabort, S. 2014. "Screening of Rice Landraces for Salinity Tolerance at Seedling Stage through Morphological and Molecular Markers." *Physiol. Mol. Biol. Plants* 20: 411-23.
- [2] Hu, S., Tao, H., Qian, Q., and Guo, L. 2012. "Genetics and Molecular Breeding for Salt-Tolerance in Rice." *Rice Genom. Genet.* 3: 38-9.
- [3] Ismail, A. M., Heuer, S., Thomson, M. J., and Wissuwa, M. 2007. "Genetic and Genomic Approaches to Develop Rice Germplasm for Problem Soils." *Plant Molecular Biology* 65: 547-70.
- [4] Thomson, M. J., de Ocampo, M., Egdane, J., Rahman, M. A., Sajise, A. G., Adorada, D. L., Tumimbang-Raiz, E., Blumwald, E., Seraj, Z. I., Singh, R. K., Gregorio, G. B., and Ismail, A. M. 2010. "Characterizing the *Saltol* Quantitative Trait Locus for Salinity Tolerance in Rice." *Rice* 3 (2-3): 148-60.
- [5] Platten, J. D., Egdane, J. A., and Ismail, A. M. 2013. "Salinity Tolerance, Na⁺ Exclusion and Allele Mining of *HKT1;5* in *Oryza sativa* and *O. glaberrima*: Many Sources, Many Genes, One Mechanism." *BMC Plant Bio.* 13: 32.
- [6] Munns, R., and Tester, M. 2008. "Mechanisms of Salinity Tolerance." *Annual Review of Plant Biology* 59: 651-81.
- [7] Apse, M. P., and Blumwald, E. 2007. "Na⁺ Transport in Plants." *FEBS Letters* 581: 2247-54.
- [8] Moradi, F., Ismail, A. M., Gregorio, G. B., and Egdane, J. A. 2003. "Salinity Tolerance of Rice during Reproductive Development and Association with Tolerance at the Seedling Stage." *Indian Journal of Plant Physiology* 8: 276-8.
- [9] Zhang, Z. H., Liu, Q., Song, H. X., Rong, X. M., and Ismail, A. M. 2012. "Responses of Different Rice (*Oryza sativa* L.) Genotypes to Salt Stress and Relation to Carbohydrate Metabolism and Chlorophyll Content." *African Journal of Agricultural Research* 7 (1): 19-27.
- [10] Yoshida, S., Forno, D. A., Cock, I. H., and Gomez, K. A. 1976. *Laboratory Manual for Physiological Studies of Rice*. International Rice Research Institute (IRRI), Manila, Philippines.
- [11] Gregorio, G. B., Senadhira, D., and Mendoza, R. D. 1997. *Screening Rice for Salinity Tolerance*. International Rice Research Institute (IRRI), Manila, Philippines.
- [12] R-STAR. 2013. *Statistical Tools for Agricultural Research*, Version 1.1. International Rice Research Institute (IRRI), Los Baños, Philippines.
- [13] Mishra, B., Singh, R. K., and Jetly, V. 1998. "Inheritance Pattern of Salinity Tolerance in Rice." *Journal of Genetics and Breeding* 52 (4): 325-31.
- [14] Peng, S., and Ismail, A. M. 2004. "Physiological Basis of Yield and Environmental Adaptation in Rice." In: *Physiology and Biotechnology Integration for Plant Breeding*, edited by Nguyen, H. T., and Blum, A. New York, N.Y. (USA): Marcel Dekker Inc., 83-140.
- [15] El-Hendawy, S. E., Hu, Y., and Schmidhalter, U. 2005. "Growth, Ion Content, Gas Exchange and Water Relations of Wheat Genotypes Differing in Salt Tolerances." *Australian Journal of Agricultural Research* 56: 123-34.
- [16] Babu, V. R. 1985. "Seed Germination, Water Uptake and Seed Reserve Utilization of Rice (*Oryza sativa* cv. Jaya) under Growth Regulator and Salinity Stressed Conditions." *Seed Research* 11: 129-35.
- [17] Fageria, N. K. 1985. "Salt Tolerance of Rice Cultivars." *Plant Soil* 88: 237-43.
- [18] Suplick-Ploense, M. R., Qian, Y. L., and Read, L. C. 2002. "Salinity Tolerance of Texas Bluegrass, Kentucky Bluegrass and Their Hybrids." *Crop Science* 42: 2025-30.
- [19] Bhowmik, S. K., Titov, S., Islam, M. M., Siddika, A., Sultana, S., and Haque, Md. S. 2009. "Phenotypic and Genotypic Screening of Rice Genotypes at Seedling Stage for Salt Tolerance." *Global Journal of Biotechnology and Biochemistry* 1 (2): 126-31.
- [20] Nakhoda, B., Leung, H., Mendioro, M. S.,

- Mohammadi-nejad, G., and Ismail, A. M. 2012. "Isolation, Characterization, and Field Evaluation of Rice (*Oryza sativa* L., Var. IR64) Mutants with Altered Responses to Salt Stress." *Field Crop Research* 127: 191-202.
- [21] Castillo, E. G., Toung, T. P., Ismail, A. M., and Inubushi, K. 2007. "Response to Salinity in Rice: Comparative Effect of Osmotic and Ionic Stresses." *Plant Production Science* 10 (2): 159-70.
- [22] Blumwald, E., Aharon, G. S., and Apse, M. P. 2000. "Sodium Transport in Plant Cells." *Biochemistry and Biophysics Acta* 1465: 140-51.
- [23] Berthomieu, P., Conéjéro, G., and Nublat, A. 2003. "Functional Analysis of *AtHKT1* in *Arabidopsis* Shows That Na⁺ Recirculation by the Phloem is Crucial for Salt Tolerance." *The EMBO Journal* 22: 2004-14.