Induced Mutation for Developing Mutant Rice Lines Tolerant to the Parasitic Weed *Striga asiatica* (L.) Kuntze

Noronirina Victorine Rakotoarisoa¹, Harimialimalala Jhonny Rabefiraisana¹, Lydia Razafinirina¹, Berthe Rasoamampionona¹, Xavier Roland Rakotonjanahary², Abdelbagi Ali Ghanim Mukhtar³, Ljupcho Jankuloski⁴ and Alice Andrianjaka¹

1. Plant Physiology Laboratory, the Department of Plant Biology and Ecology at the Faculty of Sciences-BP: 906, University of Antananarivo, Madagascar

2. Département de Recherches Rizicoles, FOFIFA, Centre National de Recherche Appliquée au Développement Rural, BP: 1690, 101-Antananarivo, Madagascar

3. Plant Breeding and Genetics Laboratory, Seibersdorf Laboratory, Joint FAO/IAEA Programme, IAEA-Laboratories, A-2444, Austria

4. Plant Breeding and Genetics Section, Joint FAO/IAEA Division of Nuclear Techniques in Food and Agriculture, Department of Nuclear Sciences and Applications, International Atomic Energy Agency, A-1400 Vienna, Austria

Abstract: This work aims to screen mutant rice lines tolerant to *Striga asiatica*. Two rainfed sensitive rice varieties B22 and F154 were used. Plants survival rates of the two parents were significantly lower respectively (9.74a and 11.83a) than those of mutant lines (55.36c to 74.36b); *Striga* plants emergence/pot were significantly higher for the parents (13.96c and 14.89c) compared to the mutants (0.12a to 1.5b); the infection rate of parents (7.37b; 7.86b) was higher compared to the mutants (2.27a to 2.74a); fertility rate/plant of parents was lower (20.98%b; 22.29%b) but much higher than mutants (72.19%b to 78.35%b); the average panicle number/plant of parents was significantly lower (0.5a; 1a) than those of mutants (1.5b to 2.4bc) and the 100 g grain weight of parents are lower (2.35a; 2.56a) than those of mutants (3.19b to 3.23b). The culture of those mutant lines may increase rice production and contribute to enhancing food security in Madagascar.

Key words: Induced mutation, mutant tolerant lines, rice (*Oryza sativa* L.), parasitic weed, *Striga asiatica* (L.) Kuntze.

1. Introduction

Rice is the staple food of the population in Madagascar. A Malagasy man eats rice on average three times a day [1] with 200 kg per inhabitant and per year, however, in terms of production, it is currently at the 19th world position with 1.2 million hectares of rice plantation [2] and an average production of 2 t/ha of paddy per year [3]. This production is insufficient, and does not cover the population needs. Rice production fails to meet the local demand. However, most of the rice cultivation in Madagascar has always been based on irrigated rice

cultivation where the available areas of which no longer respond to the population explosion in several regions of the country [4]. To ensure sustainable food security of the population, the Malagasy Government strategy encourages farmers to increase rice production through the practice of rainfed rice cultivation by using the plateaus called “Tanety” which is very large in the mid west regions of the country. But, this region has encountered a major problem which threatened the cereals plantation by the presence of a parasitic weed “*Striga asiatica*”. This witchweed plant is typically found in dry, infertile soils in semi-arid tropical grasslands and savannahs [5]. It is an annual parasitic weed, a major biotic constraint to cereal production such as maize, sorghum and rice in the middle west, north west, north

Corresponding author: N. V. Rakotoarisoa, Ph.D., lecturer researcher, research fields: plant breeding, biotechnology and genetics.

east and south west regions of Madagascar [6]. Sometimes it is perennial according to the longevity of its host [7], can stay longer more than 20 years in the soil [8] and constitutes a major threat to today's agriculture due to its minuscule tens of thousands of minute seeds produced per plant [9-11] among which up to 500,000 are produced per plant [8], its fast dissemination and its characteristics. *Striga* seeds germinate and develop "haustorium" that attaches to host roots when they receive the stimulus "Strigolactone" stimulated by host plant roots [12]. This "haustorium" in turn, tightens to the host root, draws all the required nutrients, water and growth substances from the host plant, feeds directly water, nutrients and photosynthetats through root suckers connected to the host roots system then reduces crop yields, resulting in stunting and yield reduction [13]. Once, the nutrients are enough to develop plantlet, this last emerges the soil [9, 10]. The parasitic weed *Striga asiatica* causes damage to rainfed rice culture and causes an important yield reduction from 75%-100% in Madagascar depending on soil infestation [6]. In the mid-west region where soil is much degraded and much infested, rice production varies from 0-1 t/ha [14] hence leading to food insecurity and high poverty in the country. Any rice variety was identified resistant to the parasitic plant *Striga asiatica* until now in the country and constitutes among the main problem. Mutation induction is an important complementary method of breeding crop species that speed up breeding [15]. Mutagenesis by gamma irradiation that can generate genetic variability and induce heritable change within the rice genome [15-17] was adopted for this experiment because this method has been used successfully in the genetic improvement of crop plants and has developed a lot of different cultivar such as cereals plants like rice [18, 19], ornamental or industrial plants etc., showing an important agronomical character for example earliness, high productivity, resistance to any stress [15]. This work aims to screen new rice lines tolerant to *Striga asiatica*.

2. Materials and Methods

2.1 Site of Study

2.1.1 Field

This research was conducted in the field research station of the National Research Center, FOFIFA-Kianjasoa in the middle west region of the country in Bongolava region, located at the longitude of 46°22'62.1" E; latitude of 19°03'20.3" S, and at the altitude of 889 m. The climate is sub-tropical wet of low altitude (< 500 m) to tropical wet of highland. This climate is characterized by one season of wet and raining (Nov.-April), mean temperature of 31 °C and an average rainfall of 350 mm and a second season little bit cold (May to October) with an average temperature of 25 °C and an average rainfall of 20 mm/year. The soil is characterized by net friability and sand texture, the average of carbon net is very poor, in phosphorus (< 2.5 ppm), average for carbon (1.68%) and low for nitrogen (0.25%) [20]. These edaphic conditions are favourable for *Striga asiatica* plants developments which require poor soil and dry conditions.

2.1.2 For Greenhouse Screening

Rice plantation was conducted during the summer season where climatic conditions in the screen house were: temperature ranged from 32-40 °C, luminosity: natural light; humidity: each bucket was sprayed with water twice a week during 1 month in order to ensure rice seed germination. Then, the water quantity was reduced at once a week up to pollination stage and after pollination; no water was brought into the bucket.

2.2 Plant Material Used

F154 and B22 local varieties of upland rainfed rice crops among the most appreciated by consumers and more planted by peasants, identified as sensitive varieties to *S. asiatica* in the farmer field were chosen for this experiment and received from the FOFIFA-Kianjasoa.

The dry *Striga asiatica* plants with mature capsules were collected on farmer's field one year before their use, stored in dry condition for keeping the dormancy period until 7 to 8 months.

2.3 Methods

2.3.1 Radiosensitivity Test and Plant Material Irradiation

Radiosensitivity test was performed by Razafinirina [19] at the IAEA Seibersdorf laboratories in Austria. According to her results, the two sensitive rice varieties (F-154 and B22 parents) were irradiated by gamma ray using 60Cobalt as a source at Seibersdorf laboratory in Austria in 2009. For this, 1,000 seeds per variety were irradiated at the following dose: 100, 200 Gy for F154 and 300 Gy for B22, constituting M1 seed.

2.3.2 *Striga asiatica* Pre-conditioning

Mature capsules of *Striga asiatica* were removed from plants and grinded in the mortar and then sifted in different sieve sizes up to 180 μm . Seeds were mixed with white sand such at a ratio of seeds:sands (10:90) at M2, then with 30:70 at M3. From M4, *Striga* seeds were mixed with (50:50) ratio and kept in dry condition in the Magenta until their use.

2.3.3 Field Plantation

A field of 700 m^2 was used for the experiment, ploughed and pulverized by tractor and weeding was made by manpower and then the field was divided in seven blocks of 100 m^2 each. Then, each block of 100 m^2 was subdivided in three (3) parcels of (2.5 $\text{m} \times 10 \text{ m} = 25 \text{ m}^2$) 75 m^2 which constitute 3 replications, separated by 1.5 m and 2 m between two blocks. Each block was planted by one dose of irradiated variety (M1). One seed per hole per irradiation dose per variety was planted in one parcel and the parent seeds (F154-0Gy and B22-0Gy) are planted in the middle block of those irradiated varieties. Holes were spaced by 25 $\text{cm} \times 25 \text{ cm}$. M1 seeds were planted to obtain the second generation (M2). All seeds from each plant were collected in one envelope. Two hundred (200)

rice seeds of each 500 M2 families per irradiation dose per variety were planted one by one in the same row in the field of 1 hectare (1 ha) naturally infested by *S. asiatica*. Rice plants were spaced by 25 $\text{cm} \times 25 \text{ cm}$ and each block was spaced by 2 m and 1 m for each parcel. Since M2 generation, parent seeds were planted in the same parcel as the irradiated variety in the 3 middle rows of the parcel in order to facilitate the observation of the phenotypic differences between plants.

2.3.3.1 Soil Infestation

The soil is naturally infested by *S. sciatica* but seeds repartitions in soil are not homogenous. From M2 generation, a selection pressure was practiced. The preconditioned *Striga* seeds were used for artificial inoculation in order to ensure the presence and well repartition of the parasitic plant in the soil, also to ensure interaction between host plant roots and the parasitic. For this, one (1) pinch around of 3,000 of preconditioned *S. asiatica* seeds was inoculated and scattered per hole. No fertilizer was brought to the field. These conditions promote *S. asiatica* plants attack to the host plant. The dose rate of the *Striga* seeds was increased per year.

2.3.4 Greenhouse Screening

From M4 to M6 generations, 5 rice seeds per pot from the same plants, the same dose of irradiation and from the same variety of the selected lines were planted in the pot of 10 L containing soil. Severe artificial infestation was done in each pot by inoculating and scattering *S. asiatica* seeds in two (2) levels of 8 cm and 12 cm soil depth. Two pinches around of 5,000 *S. asiatica* seeds were inoculated in each level then, covered by soil in order to ensure the interaction between host plant roots and the parasitic weeds. No fertilizer was applied during greenhouse experiments.

2.3.5 Criteria and Selection Method

Mass selection was applied at M2 generation by considering the plant having the highest full grain and fewer symptoms on leaves, followed by "individual

selection” by “Pedigree screening” since M3 to M6. All parameters were assessed to determine the impact of *S. asiatica* attack to the advanced selected lines M5 and M6 plants generation. For the infection rate, the total number of infected leaves per plant was recorded and assessed according to Haussmann, et al. [21]. The tolerance of the selected plant was determined by combining the infection rate per plant, the plant height, the *Striga* plants emerging soil per pot with the seeds fertility number per plant.

2.3.6 Parameters Considered

The plant height, *Striga asiatica* plants emerged number/pot (counted at 80 days after sowing), host plant survival rate, infection rate per plant were assessed to determine the impact of the *S. asiatica* attack to the host plant tolerance and the 100 seeds weight (g) per genotype, the fertile/sterile seeds number per plant, the average panicle number per plant were assessed to determine the agronomic impact of the *S. asiatica* attack to host plant.

2.4 Statistical Analysis

Data were collected and subjected to ANOVA (Analysis of Variance) analysis using the software R version 3.4.1 followed by a comparison of means using Tukey HSD (Honestly Significant Difference) test at the significance level of $p \leq 0.05$. Interaction or relationships between parameters were assessed by the ACP (Principal Component Analysis) with the Pearson correlation test.

The Fig. 1 shows the method adopted to develop morphologically mutant tolerant rice lines.

3. Results

3.1 The Number of Selected Genotypes at Different Stage M1, M2, M3, M4, M5 and M6

According to the characters required at M2 for mass screening, all plants having high spikelet fertility and less symptoms on leaves were collected for the next screening. The total number of the different selected genotypes (families) at M2, M3, M4, M5 and M6

generations is given in Table 1.

Table 1 shows that at M2 generation, 500 families per irradiation dose and per variety were fixed and chosen to continue the next step of screening, those which have the maximum of seeds fertility and fewer symptoms on leaves. This number was fixed in order to ensure the probability of getting 10^{-3} mutants by screening more genotypes 100,000 according to Van Harten [16]. And at M3 and M4, the number of selected genotypes decreased per irradiation dose and per variety, because after increasing the dose of infestation in each hole “the ratio [*Striga* seeds/sands]” from 20/80 at M2; 30/70 at M3; and 50/50 at M4), most of the susceptible and sensitive genotypes died and the survived plants were selected for the next screening season. The more the dose rate of the *S. asiatica* seeds inoculated in the hole was increased, the less the putative tolerant lines number selected decreased. This result demonstrates the effectiveness of the high density of *Striga* seeds inoculated in the hole which were very close to the host roots plants [11] and then promoted interactions between parasitic and the host plant roots. The difference of the selected genotypes numbers at M5 and M6 generations are not significant. The *S. asiatica* ratio was kept at 50:50 and the total number of surviving plants at M6 generation did not change anymore. This situation means that the genotypes characters are almost stable at M6. This result confirms that observed by Van Harten [16] who mentioned that the characters of cereal plant issue of induced mutation by irradiation is almost stable at M6 generation with the autogamous plant, case of rice in our experiment.

3.2 Impact of *Striga asiatica* Attack to the Physiological Parameters of the Host Plant

Table 2 shows the impact of *Striga asiatica* attack to the host plant of all selected genotypes at M6 generation, planted in the greenhouse under severe infestation by considering different parameters. Table 2 shows the average number observed within the

Fig. 1 Scheme of Development method of mutant rice lines B22 and F154 through conventional mutagenesis developed by Rakotoarisoa, N. V., et al [22].

Table 1 Result of the total number of the selected rice genotypes per generation.

Genotype	Nb of selected families at M2	Nb of selected genotypes at M3	Nb of selected genotypes at M4	Nb of selected genotypes at M5	Nb of selected genotypes at M6
F154-100 Gy	500	180	15	8	6
F154-200 Gy	500	171	8	5	4
B22-100 Gy	500	195	15	9	7
B22-200 Gy	500	188	14	8	6
B22-300 Gy	500	145	20	5	4

Nb: number.

Table 2 Impact of *S. asiatica* attack to the rice host plant at M6 generation in the greenhouse.

Genotype	<i>S. asiatica</i> plant emerged number/pot	Average infection rate/plant (host damage)	Survival rate (%)	Plant height (cm)
F154 0 Gy	13.96b	7.37b	11.83a	63.62a
F154 100 Gy	0.12a	2.27a	73.35b	69.7b
B22 0 Gy	14.89b	7.86b	9.74a	46.5b
B22 100 Gy	0.5a	2.2a	74.36b	42.7a
B22 200 Gy	0.7a	2.3a	55.36b	49.5c
B22 300 Gy	1.5a	2.74a	74.25d	52.8d

Different letters mean significant statistical differences between treatments at $p \leq 0.05$ values within columns (issues for the same variety), the same letters are not significantly different.

same selected genotypes from the same variety and the same dose of irradiation.

3.2.1 Regarding the Number of Emerged *S. asiatica* Plant per Pot

The results show that the average *Striga asiatica* plants emerged aboveground in each pot for the parent varieties F154 0 Gy and B22 0 Gy respectively 13.96b, 14.89b are much higher than those of the selected irradiated M6 genotypes 0.12a to 1.5a. Differences are significant at $p \leq 0.05$. For the advanced lines (issues of 100, 200, 300 Gy) of B22, the number of *Striga asiatica* plants emerging soil was increased with the dose of irradiation but there is no significant difference between numbers from different genotypes. Among the advanced selected lines, B22-300 Gy have the highest *S. asiatica* plants emerged per pot, which means that the high dose of 300 Gy has increased the roots stimulation of Strigolactone within the host plant B22 irradiated at 300 Gy, but this value 1.5a is significantly lower than those of the parent varieties B22-0 Gy (14.89b) and F154-0 Gy (13.96b). These results demonstrate that the irradiation using gamma ray has led to the reduction of the Strigolactone amount stimulated by host plants roots which limits the *Striga asiatica* seeds germination then the low and few *Striga asiatica* plants emergences. This low induction of *Striga* seeds germination constitutes a major component of resistance in these lines as observed [23] by working on Faba bean against *Orobanche foetida*. This situation needs to confirm by Strigolactone contents analysis of those improved lines. Hence, irradiation using gamma ray at 100-200

and 300 Gy has improved host plant rice genome leading to its tolerance to *S. asiatica* attack.

3.2.2 For the Infection Rate per Plant

All of the advanced selected genotypes have very low infection rate per host plant varying from 2.20a to 2.74a than those of the parent varieties F154 and B22 respectively 7.37b to 7.86b. Differences between parents are not significant, but significant between advanced lines and parents. The infection rates 7.37b to 7.86b of these parents correspond to “sensitive” variety [21], which are their right situation during our farmer’s field surveys and prospecting on 2008. There is no difference also between selected genotypes which exhibit the stability of their characters in front of *S. asiatica* attack. These situations determine that all selected plants are ranked as “tolerant to resistant” genotypes to the parasitic weeds according to Haussmann, et al. [21].

This parameter is linked to the number of *S. asiatica* plants emerged aboveground, depending also on the density and the quality and the repartition of *Striga* seeds in the holes. When *Striga* seeds do not reach the required dormancy period, and if the seeds quality is too bad, they cannot germinate even if they receive stimuli from host plants. The emergence of *Striga* plants in the parent variety indicates that *Striga* seeds have successfully lifted dormancy. Because when good quality of *S. asiatica* seeds underground surrounding host roots plants received stimuli from host plant, they germinated [9, 10] and turn to the host plant roots by positive chemotropism, attached to the host roots via “haustorium” and penetrate into its

xylem, then divert nutrients to their benefit [12]. When nutrient extracted from host plant is enough, *S. asiatica* plant emerges aboveground. This number of *S. asiatica* plants emerging soil should reflect the number of underground *Striga* germinated and closely linked to the infection rate on leaves of host plant. The more the infection rate, the more the tested variety has high density of emerged *Striga* plants. The low *Striga* plants emergence could be explained by the low *Striga* seeds germinated due to the lack or very low stimuli from host roots plants. The ability of the host plant to avoid penetrating of haustorium into its xylem is genotype dependant. By using the same dose ratio of (*Striga* seeds:sands) in the pot, the parent varieties F154-0 Gy and B22-0 Gy have shown high infection rate respectively 7.37b and 7.86b than the selected advanced M6 genotypes, which varies from 2.2a to 2.74a. According to these results, the plant genome of the M6 genotypes may possess such tolerance or resistance barrier within the xylem or endoderm or pericycle layers, because the infection rate visible on leaves is linked totally to the presence or absence of the interaction between host plants roots and the parasite *Striga asiatica* infection. For this, all selected advanced genotypes are classified as mutant tolerants to *S. asiatica* with very low infection rate on leaves from 2.2a to 2.74a.

3.2.3 About the Survival Rate of Host Plants

Under severe infestation of *Striga* seeds/sands ratio (50/50) at two levels 8 and 12 cm soil depth, the survival rate of host plants shows that the parent varieties F154-0Gy and B22-0Gy cannot support the *Striga asiatica* attack, respectively 11.83%a and 9.74%a only are surviving. These percentages are significantly lower than those of the advanced selected genotypes having more than 68.4%b up to 80.5%b surviving plants. For B22 variety, the survival rate increases with the dose of irradiation, except at 200 Gy. This result explains that B22 variety supports well the high dose of irradiation 300 Gy which leads to the improvement of its tolerance character to the parasitic

plant *S. asiatica* by keeping 82.3% of surviving plants under severe infestation of *S. asiatica*. For F154 variety, the dose of 100 Gy is enough to improve its rice genome character and can survive 73.35b plants at M6. These survival rates explain that all parent varieties are more sensitive to the attack of *S. asiatica* than the advanced lines because after their germination, when the parasite attacks the host plant during vegetative phase, plants are wilting and die when susceptible, but can survive if host plant is sensitive but almost sterile or with high sterility [24] (photos 1 and 3). The case is different for the selected genotypes, their survival rates are significantly superior to those of parent varieties. These results confirm that the selected genotypes may possess different resistance or tolerance mechanism barrier to avoid *S. asiatica* penetrating and attack. This situation needs to confirm also to determine this kind of tolerance or resistance within the plant genome.

3.2.4 About the Plant Height

The parent variety F154 is shorter than the advanced selected lines M6-F154-100 Gy with $p \leq 0.05$. But for B22, the genotype B22-100 Gy is the shortest stature 42.7a, followed by the parent variety 46.5b then, the selected lines B22-200 Gy and 300 Gy (49.5c and 52.8d). For the B22 improved lines, the plant height increased when irradiation dose increased. The short stature of B22-100 Gy could not be due to *Striga* infection nor its attack, because both its infection rate and *Striga* plants emergence per pot are significantly very low (0.2a and 2.2a) than that of the parent variety. The differences of plant height between selected advanced genotypes are significant at $p \leq 0.05$. So, this difference of plant height among the improved lines can be due to irradiation effect. This last can reduce plant height [15, 18, 19].

These results mean that all selected genotypes are classified as mutant genotypes tolerant to *Striga asiatica*, result similar to that of Haussmann, et al. [21].

3.3 Assessing the Impact of *Striga asiatica* Seeds Inoculation in Each Hill on the Agronomical Characters of M6 Genotypes

3.3.1 Panicle Fertility Rate

Table 3 below describes that both parent varieties F154-0 Gy and B22-0 Gy have the average lowest panicle fertility rate respectively 21.07%a and 22.46%a compared to those of advanced selected genotypes from F154-100 Gy and B22-100-200 and 300 Gy which are respectively 85.71%c, 86.87%b, 83.33%b and 85.02%b. No difference was observed between selected genotypes of B22. Moreover, the more the irradiation dose increases, the more the panicle fertility rate increases. In general, all selected genotypes have a fertility rate above 83% under severe infestation at two levels of 8 and 12 cm. This character proves that all selected genotypes are “mutants tolerant to *S. asiatica*” because under severe infestation, the host plant (not improved) dies according to Andrianaivo, et al. [6], or in authors’ experiment it dies or becomes highly sterile if it can survive.

In that case plant is considered as susceptible when after 1 or 2 weeks of germination, there is underground attack and the host plant dies. It is scored as sensitive if it can survive after *S. asiatica* attack showing high infection symptoms on leaves after pollination period or during reproductive phase. The Pearson’s correlation is positively high correlated with $r = 0.820^{***}$ $p < 0.001$ for F154 and 0.756^{***} with $p < 0.001$ for B22. Grain yield losses due to *S. asiatica* parasitism can attain 100% in sensitive cultivars under high infestation levels, particularly under drought condition; this result confirms those observed by Bettina, et al. [25].

3.3.2 The Average Panicle Number per Plant

For F154, the panicle number per plant of the improved line F154-100 Gy is significantly superior (1.03b) than that of the parent (0.5a). The same case was observed for the B22 genotypes, all selected lines

have shown high panicle number 1.75b to 3.0c than the parent variety B22-0Gy. Differences are significant between lines from irradiation doses. The more the irradiation dose increases, the more the panicle number per plant increases. To describe their tolerance to the parasitic weed *S. asiatica*, authors have to consider the fertile and sterile seeds number produced per panicle.

3.3.3 The 100 Seeds Weight

For the two parent varieties F154-0 Gy and B22-0 Gy, their 100 seeds weight (g) in Table 3 are different 2.35a for F154-0Gy and 2.56a for B22-0Gy, but the difference is not significant. When compared to the selected advanced lines, their 100 seeds weights are significantly lower than those of selected advanced lines (3.19b to 3.23b). However, no significant difference is observed between 100 seeds weight of the selected advanced tolerant genotypes.

3.3.4 Fertility Rate

Considering the methods adopted during plantation that no fertilizer was brought to ensure that soil is poor in order to promote *S. asiatica* attack to the host plant and the presence of interaction between roots, the fertility rate of advanced selected lines has high fertility rate. These characters were improved also through the rice genotypes irradiation and lead to phenotypic differences. By considering the average fertile seeds number per plant, in reality, some rice genotypes are 100% fertility (without sterile seed) and no *S. asiatica* plants emerging soil, those lines are considered as “mutant resistant to *S. asiatica*”.

3.4 Analysis of Relationship between Parameters

Table 4 shows Pearson’s correlation coefficients (r) describing association of different parameters using F154-variety.

The PCA (Fig 2) observation has shown 98.99% of the information according to the axes F1 & F2. For the correlation study on this part, the symbols * and *** represent the significant levels of $p < 0.05$ and $p < 0.01$, respectively.

Table 3 Impact of *S. asiatica* attack to the agronomical characters of the M6 rice mutant host plants.

Genotype	Average Fertile seeds number/plant	Average sterile seeds number/plant	Panicle Fertilty rate (%)	Average panicle number/plant	100 seeds weight (g)
F154 0 Gy	5.5a	20.6b	21.07a	0.5a	2.35a
F154 100 Gy	22.8b	3.8a	85.71c	1.03b	3.19b
B22 0 Gy	5.1a	17.6b	22.46a	1.00a	2.56a
B22 100 Gy	19.8b	3.8a	86.87b	1.75b	3.23b
B22 200 Gy	20.0b	4.0a	83.33b	1.96b	3.19b
B22 300 Gy	21.0b	3.7a	85.02b	3.0c	3.20b

Different letters mean significant statistical differences between treatments at $p \leq 0.05$, values within columns with the same letter are not significantly different.

Table 4 Pearson's correlation coefficients (r) describing association of different parameters using B22 and F154-varieties.

Pearson Correlation			B22		F154			
			Pearson's r	p	Pearson's r		p	
Fertile_seeds_number	-	Weight_100seeds	0.527	***	< .001	0.734	***	< .001
Fertile_seeds_number	-	Height	0.651	***	< .001	0.404		0.077
Fertile_seeds_number	-	Sterile_seeds_number	-0.756	***	< .001	-0.877	***	< .001
Fertile_seeds_number	-	Striga_number	-0.737	***	< .001	-0.841	***	< .001
Fertile_seeds_number	-	Haussman_score	-0.832	***	< .001	-0.893	***	< .001
Weight_100seeds	-	Height	0.350	*	0.027	0.225		0.341
Weight_100seeds	-	Sterile_seeds_number	-0.638	***	< .001	-0.803	***	< .001
Weight_100seeds	-	Striga_number	-0.565	***	< .001	-0.742	***	< .001
Weight_100seeds	-	Haussman_score	-0.556	***	< .001	-0.752	***	< .001
Height	-	Sterile_seeds_number	-0.228		0.158	-0.489	*	0.029
Height	-	Striga_number	-0.231		0.151	-0.545	*	0.013
Height	-	Haussman_score	-0.231		0.152	-0.175		0.460
Sterile_seeds_number	-	Striga_number	0.801	***	< .001	0.825	***	< .001
Sterile_seeds_number	-	Haussman_score	0.893	***	< .001	0.820	***	< .001
Striga_number	-	Haussman_score	0.871	***	< .001	0.731	***	< .001

* and *** represent the significant levels of $p < 0.05$ and $p < 0.01$, respectively.

During the test of all advanced genotypes of F154 and B22, the infection rates of host plant vary from 2.2a to 7.86b, this score is very low, and this parameter is *strongly correlated negatively* with their 100 seeds weight with $r = -0.752^{***}$ and with fertile seeds number $r = -0.893^{***}$. This situation means that there is high negative correlation between these parameters. The more the infection rate increases, the less both 100 seeds weight and fertile seeds number increase. However, this damage score is *strongly correlated positively* with the *Striga* plants emerged number for F154, because the *Striga* number is so high with parent varieties with $r = 0.731^{***}$ and very low for the selected advanced genotypes. These situations confirm the tolerance of the selected

genotypes to the attack of *S. asiatica*.

The infection rates of the parent varieties F154-0Gy and B22- 0Gy are in positive highly correlated with high sterile seeds number and the high *Striga* plants emerging soil with $r = 0.825^{***}$; these parent varieties parameters are in *strongly negatively correlated* with their plants height. The more the *Striga* plants emerged increase, the more the plant height decreases. All of these parameters are in negative correlation to those of their advanced selected genotypes. These last (B22 100 Gy, B22 200 Gy, B22 300 Gy and F154 100Gy) share common characteristics in positive high correlation with high survival rate, high fertile seeds number, low *Striga asiatica* plants emerging soil, high 100 seeds weight

PCA result:

Fig. 2 Observation of Principal component analysis (PCA).

Striga No: *Striga* plants number; Fertile seeds No: Rice fertile seeds number; Sterile seeds No: rice sterile seeds number.

and low or without infection symptom rate. These parameters of advanced selected genotypes are in opposite of that mentioned by Graves [26] who reported that the infection inevitably results in yield and quality of crops and host damage becomes noticeable even before the parasite emerges from the soil.

The Fig. 2 below is showing the correlation between parameters and treatment factors.

4. Conclusion

Physical treatments by gamma ray were used, followed by individual selection for heritable changes of specific genotypes, and irradiation has led to positive impact on agronomical characters of the tested selected genotypes by increasing the number of filled grain and 100 seeds weight, reducing the empty seed number and the number of *S. asiatica* plants emerged per pot. The irradiation of the explant at 100-200 and 300 Gy doses has improved the genome within the selected mutant rice lines by improving their tolerance to the attack of *S. asiatica*, reducing the

damage score of the tested genotypes under severe infestation. Mutagenesis has been used as a potent method of enhancing variability for crop improvement; hence, 27 were identified as tolerant to the parasitic weed *Striga asiatica*. We envisage that these selected mutant lines will be multiplied and will be tested on farmer's field and mutant will be identified by molecular characterization. The plantation of these

Photo 1 B22-0Gy plants attacked by *S. asiatica* with more *Striga* plants emerged, stunting and sterile totally. Sensitive variety.

Photo 2 Selected B22-300 Gy M6, tolerant, fertile, no symptom of attack and without *Striga* plant emerged under severe infestation.

Photo 3 F154-0Gy parent variety, with high symptom attack on leaves, high damage score, totally sterile, dried and died, classified as sensitive variety.

Photo 4 F154-100 Gy, selected mutant resistant lines, without symptom attack on leaves, no *Striga* plant emerged, high fertility rate under severe infestation.

mutant lines may increase rice production on farmer level then in the country, and can contribute to the food security of the population.

Acknowledgements

Authors are very thankful and grateful to the IAEA (International Atomic Energy Agency, Joint FAO/IAEA Division of Nuclear Techniques in Food and Agriculture in Austria), for their supporting and collaboration of this project.

Authors do not forget also the FOFIFA team with the head of the station, technicians and manpower for their kind collaboration. Authors thank the anonymous reviewers for comments that greatly improved the manuscript.

References

- [1] Accessed in 2017. <https://www.madagascar-vision.com/riz-culture-madagascar/>.
- [2] Website of Production de riz à Madagascar. <http://www.dggfpe.mg/index.php/2017/06/23/production-de-riz-madagascar-19eme-rang/>.
- [3] Maep. 2005. "Ministère de l'Agriculture, de l'élevage et de la Pêche." *Filières de l'Agriculture, de l'Elevage et de la Pêche, et Actions du Ministère de l'Agriculture, de l'Elevage et de la Pêche*. Filière maïs fiche, No.108, p. 10. (in French)
- [4] Spallek, T. M., Mutuku, J., and Shirasu, K. 2013. "The Genus *Striga*: A Witch Profile." *Molecular Plant Pathology* 3 (9): 1-9.
- [5] Husson, O., Michellon, R., Charpentier, H., Razanamparany, C., Moussa, N., Naudin, K., et al. 2008. "Le contrôle du *Striga* par les systèmes SCV (Semis direct sur couverture végétal permanente)." In *Manuel pratique du semis direct à Madagascar*, Volume I, Chapitre 3, §3.1, p. 20. (in French)
- [6] Andrianaivo, A. P., Kachelriess, S., Kroschel, J., and Zehrer, W. 1998. *Biologie et gestion de Striga à Madagascar*. FOFIFA, DPV, GTZ, p. 65. (in Corsican)
- [7] Sallé, G., Raynal-Roques, A., Tuquet, C., Andary, C., Dembele, B., and Diallo, S. 1994. "Striga Research in Mali, Burkina Faso and Senegal." In *Biology and Management of Orobanchae. Proceedings of the Third International Workshop on Orobanchae and Related Striga Research*, edited by Pieterse A. H., Verkleij, J. A. C., and Borg, S. J., Amsterdam, pp. 700-9.
- [8] Doggett, H. 1988. *Sorghum*, 2nd ed. Essex, England: Longman Scientific & Technical.

- [9] Jackson, M. B., and Parker, C. 1991. "Induction of Germination by a Strigol Analogue Requires Ethylene Action in *Striga hermonthica* but Not in *S. forbesii*." *J Plant Physiol* 138: 383-6.
- [10] Logan, D. C., and Stewart, G. R. 1991. "Role of Ethylene in the Germination of the Hemiparasite *Striga hermonthica*." *Plant Physiology* 97: 1435-8.
- [11] Cechin, I., and Press, M. C. 1993. "Nitrogen Relations of the Sorghum-*Striga hermonthica* Host-Parasite Association: Germination, Attachment and Early Growth." *New Phytol.* 124: 681-7.
- [12] Bouwmeester, H., Matusova, R., Zhongkui, S., and Beale, M. H. 2003. "Secondary Metabolite Signaling in Host-Parasitic Plant Interactions." *Plant Biology* 6: 358-64.
- [13] Nail, K., Kriticos, D. J., Scott, J. K., Yonow, T., and Ota, N. 2014. "*Striga asiatica*." *Harvest Choice Pest Geography* 2: 1-6.
- [14] Rakotondramanana. 2004. "Riziculture et SCV." *Dans Filière de l'Agriculture, de l'Élevage et de la Pêche*, et Actions du Ministère de l'Agriculture de l'Élevage et de la Pêche. Fiche No. 108, p. 5. (in French)
- [15] Fao/laea. 2018. "Manual on Mutation Breeding." In *Plant Breeding and Genetics Sub-programme*, 3rd ed., edited by Spencer-Lopes, Forster, B. P., and Jankuloski, L., Joint FAO/IAEA Division of Nuclear Techniques in Food and Agriculture, Vienna-Austria, p. 399.
- [16] Van Harten, A. M. 1998. *Mutation Breeding, Theory and Practical Applications*. Sweden: Svalof Weibull, Plant Breeding Company, p. 302.
- [17] Mba, C. R. 2013. "Induced Mutations Unleash the Potentials of Plant Genetic Resources for Food and Agriculture." *Agronomy* 3: 200-31. doi:10.3390/agronomy3010200. ISSN: 2073-4395.
- [18] Rakotoarisoa, N. V. 2008. "Développement de lignées productives et de lignées tolérantes au froid par la mutagenèse in vitro à partir des variétés de riz (*Oryza sativa* L.) Rojofotsy, Malady et IR58614." PhD thesis, Université d'Antananarivo-Madagascar. (in French)
- [19] Razafinirina, L. 2011. "Test de radio sensibilité de quelques variétés de riz pluvial et de maïs cultivées à Madagascar." Mémoire pour l'obtention du diplôme d'Etudes Approfondies en Biologie et Ecologie Végétales, Parcours Physiologie et Biotechnologie Végétales, Faculté des Sciences, Université d'Antananarivo-Madagascar, p. 44. (in French)
- [20] Razafimahefa, P. 2010. "Contribution à l'étude des paramètres physico-chimiques des sols du Moyen-ouest et Moyen-est de Madagascar dans le cadre de la valorisation de la culture du riz." Mémoire de DEA en chimie minérale, Université d'Antananarivo-Madagascar, p. 110. (in French)
- [21] Haussmann, B. I. G., Hess, D. E., Welz, H. G., and Geiger, H. H. 2008. *Erratum to Improved Methodologies for Breeding Striga-Resistant Sorghum*. University of Hohenheim, Institute of Plant Breeding, Seed Science and Population Genetics, pp. 195-211.
- [22] Rakotoarisoa, N. V., Rabefiraisana, J. H., Rasoamampionona, B., Tokiniaina Fandresena, M. E., Razafindrasoa, M. A., Rakotonjanahary, X. R. T., and Andrianjaka, A. 2017. "Sélection des géotypes de maïs tolérants à *Striga asiatica* à Madagascar." *Dans Actes du Forum de la Recherche à Fianarantsoa*, 29-30 Novembre 2017 à Fianarantsoa-Agrobiodiversité, organisé par le Ministère de l'Enseignement Supérieur et de la Recherche Scientifique, pp. 257-62. (in Corsican)
- [23] Mejrik, S., Yassine, M., Omrane, B., and Mouldi, S. 2018. "Orobanche Foetida Resistance in Two New Faba Bean Genotypes Produced by Radiation Mutagenesis." *International Journal of Radiation Biology* 3002: 1362-3095.
- [24] Accessed in 2017. <http://www.amdp.mg/index.php/2017/10/31/riziculture/>.
- [25] Bettina, B. I. G., Haussmann, D. E., Hess, B. H., Günter, W., and Geigera, H. H. 2000. *Improved Methodologies for Breeding Striga-Resistant Sorghums*. University of Hohenheim, Institute of Plant Breeding, Seed Science and Population Genetics, 70593 Stuttgart, Germany International Crops Research Institute for the Semi-Arid Tropics, BP 320, Bamako, Mali, p. 17.
- [26] Graves, J. D. 1995. "Host-Plant Responses to Parasitism." In *Parasitic Plants*, edited by Press, M. C., and Graves, J. D. London: Chapman & Hall, pp. 206-25.